

Full Length Research Paper

# Specific detection of *Aspergillus fumigatus* in sputum sample of pulmonary tuberculosis patients by two-step PCR

Sunita Bansod<sup>1\*</sup>, Inderchand Gupta<sup>2</sup> and Mahendra Rai<sup>2</sup>

<sup>1</sup>Biotechnology Division, Defense R and D establishment, Jhansi Road, Gwalior-474002, India.

<sup>2</sup>Biotechnology Department, Amravati University, Amravati-444107, India.

Accepted 14 November, 2020

The incidence of life-threatening systemic fungal infections has been increasing in recent years, and the increasing incidence has been correlated with increasing numbers of immunocompromised patients. We developed a two-step PCR assay that specifically amplifies a region of the 18S rRNA gene that is highly conserved in *Aspergillus fumigatus*. This assay allows direct and rapid detection of down to 10 fg of *Aspergillus fumigatus* DNA corresponding to 1 to 5 colony forming unit (CFU) per ml of sputum sample of pulmonary tuberculosis patients. The two primer pairs that produced PCR products with the highest sensitivity and species specificity were the *Aspergillus fumigatus* primers AFU7S and AFU7AS, which amplified a fragment of 405 bp, followed by AFU5S and AFU5AS, which produced an internal fragment of 236 bp. PCR has been shown to be a highly sensitive diagnostic tool for the detection of infectious fungi specimens. Our point to the considerable potential clinical value of this simple, specific, rapid, and inexpensive PCR assay is for improving the means of early diagnosis of systemic aspergillosis tuberculosis patients.

**Key words:** *Aspergillus fumigatus*, Polymerase chain reaction (PCR), pulmonary tuberculosis.

## INTRODUCTION

The incidence of life-threatening systemic fungal infections has been increasing in recent years, and the increasing incidence has been correlated with increasing numbers of immunocompromised patients (Anaissie et al., 1989; Bodey et al., 1992; Beck-Sague, and Jarvis, 1993; Groll et al., 1996; Denning et al., 1998; Kami et al., 2000). Patients at the greatest risk are those with prolonged periods of pulmonary tuberculosis after intensive immunosuppressive chemotherapy, for example, during treatment for acute leukemia or after bone marrow transplantation. Particularly on the increase are invasive infections with *Aspergillus fumigatus*, resulting in high mortality rates or, if the patient survives, causing high levels of morbidity that often limit further antileukemic therapies. Antifungal prophylaxis is controversial and is not generally practiced (Arning et al., 1994; Cafferkey et

al., 1994; Chandrasekar et al., 1995; Wald et al., 1997; Lin et al., 2001). Moreover, amphotericin B is toxic. In contrast to other infections, only limited conventional diagnostic tools with poor sensitivity and reliability are available for early detection of invasive aspergillosis (Walsh et al., 1991; Denning et al., 1994; Guiot et al., 1994; Hughes et al., 2002.) with the systemic infection frequently being diagnosed late or confirmed only at autopsy (Bodey et al., 1992; Pfaffenbach et al., 1994; Chandrasekar et al., 1995; Groll et al., 1996). The PCR test commonly used in the initial period of infection during pulmonary tuberculosis are often insufficient not only for accurate early diagnosis but also for monitoring of the subsequent course of invasive aspergillosis (Anaissie et al., 1989; Blum et al., 1994; Guiot et al., 1994; Logan et al., 1994; Obayashi et al., 1994; Hearn et al., 1995; Styne et al., 1995; Verweij et al., 1995; Walsh et al., 1995a, b; Kappe et al., 1996; Caillot et al., 1997; Swanink et al., 1997; Lewin et al., 1999; Loeffler et al., 2000a; Kearns et al., 2001; Palladino et al., 2001). In view of the low specificity and sensitivity rates of these methods, the

\*Corresponding author. E-mail: [sunita.bansod@rediffmail.com](mailto:sunita.bansod@rediffmail.com)

diagnosis of an invasive aspergillosis can be proven conclusively only by positive histology or culture results (Bart-Delabesse et al., 2001). However, establishing cultures from blood and bronchoalveolar lavage (BAL) samples is often unsuccessful due to the low yields of CFU, and in the case of immunocompromised high-risk patients who are febrile, pulmonary tuberculosis, neutropenic, thrombocytopenic, and often seriously ill, tissue biopsy specimens, in general, are not available.

Many authors have reported detection of *Aspergillus* nucleic acids by PCR for improved diagnosis of invasive aspergillosis. Target sequences, either panfungal ribosomal DNA (rDNA) (Einsele et al., 1997; Hebart et al., 2000; Hendolin et al., 2000; Lass-Flörl et al., 2001; Loeffler et al., 2000; Loeffler et al., 1998) or *Aspergillus*-specific mitochondrial

DNA (mtDNA) (Bretagne et al., 1998; Costa et al., 2002; Jones et al., 1998) or rDNA sequences (Buchheidt et al., 2002; Buchheidt et al., 2001; Kami et al., 2001; Skladny et al., 1999) have been amplified from BAL fluids (Skladny et al., 1999; Buchheidt et al., 2002) serum (Bretagne et al., 2002; Costa et al., 2002), or whole blood (Buchheidt et al., 2002; Hebart et al., 2000; Einsele et al., 1997; Lass-Flörl et al., 2001). The diagnosis of mycotic infections in pulmonary tuberculosis patients by molecular methods such as Southern blot analysis has been performed successfully with lung. More sensitive and rapid detection assays have been established by use of the PCR method, particularly following the identification and sequencing of multicopy gene templates in a range of fungi and other organisms. PCR assays for the detection of fungal nucleic acids may be the optimal diagnostic approach because they are potentially more sensitive than current culture-based methods and may be designed to encompass the desired range of genera and specimen types. Previous studies evaluating PCR-mediated detection of *Aspergillus* species showed significantly improved sensitivity but involved assays with different methods and objectives, partly to optimize culture assays (Burg et al., 1989; Makimura et al., 1994; Hayden et al., 2001.) and partly for typing in epidemiological studies (Aufavre et al., 1992; Girardin et al., 1994; Birch et al., 1995; Groll et al., 1996). Therefore, the results of different groups are not consistent or comparable. By using PCR primers specific for the multicopy 18S rRNA gene, very small amounts (down to 1 pg) of genomic DNA from *A. fumigatus* have been detected, the sensitivity being increased to 100 fg by (Spreadbury et al., 1993).

Melchers et al. (1994) first described a PCR assay for the detection of DNA from an *Aspergillus* sp. in immunocompromised patients with primers based on the coding sequence of the 18S rRNA gene which is highly conserved and which is amplified some hundredfold in the *Aspergillus* genome. PCR products were obtained from BAL samples of immunocompromised while no amplicons were obtained from immunocompetent individuals. However, Yamakami et al. (1996) first described the use of a two-step PCR to detect *Aspergillus* spp. in blood with increased spe-

specificity and sensitivity, but that study was performed with only a small number of patients. Einsele et al. (1997) described a PCR assay with subsequent Southern blot analysis, which allowed the detection of fungal pathogens (including *Aspergillus* spp.) in blood samples.

In order to achieve an improved, specific, and rapid means of detection of *Aspergillus fumigatus* in clinical specimens, we developed a two-step PCR assay for sputum samples. Two optimal pairs of oligonucleotide primers derived from sequences of the 18S rRNA gene, which are specific for *A. fumigatus*. The assay was evaluated for its sensitivity and specificity *in vitro* and was used to analyze clinical samples of immunocompromised patients.

## MATERIALS AND METHOD

### Clinical samples and isolation

*A. fumigatus* test strains were isolated from the sputum samples of pulmonary tuberculosis patients from the civil hospital of Washim, Washim District, Maharashtra state, India. Sputum samples (2 - 4 ml) were obtained under sterile conditions. The samples were inoculated in appropriate sterile Czapek Dox Agar plates at 25°C for 3 to 4 days. The cell density of the fungal suspensions (conidia) was determined microscopically by counting the cell number in a Neubauer cell chamber.

### DNA isolation

DNA was isolated from larger scale (25 ml) cultures. Mycelium were separated by filtration and washed three times in cold buffer (20mM sodium citrate pH 5.8 in 1 M sorbitol), ground in liquid nitrogen and resuspended in lyses buffer (50 mM Tris-HCl (pH 8), 50 mM EDTA, 3% sodium dodecyl sulfate, and 1%  $\beta$ -mercaptoethanol). The suspension was then incubated at 65°C for 1 h. After the cell debris had been pelleted, the lysate was extracted once with phenol-chloroform (1:1) and once with a chloroform-isoamyl alcohol (24:1). The DNA was precipitated by the addition of 0.2 volume of 3 M potassium acetate and 0.5 volume of isopropanol; after centrifugation, the DNA pellets were washed with 70% ethanol and resuspended in buffer (10 mM Tris-HCl pH-8, 1mM EDTA).


The purification of fungal DNA was performed by conventional extraction (Sambrook et al., 1989). The DNA was precipitated by the addition of 0.7 volume of isopropanol, pelleted, and washed once with 70% ethanol and air-dried. The DNA concentration was assessed by spectrophotometer at 260 and 280 nm.

### Oligonucleotide primers, primer sequences, and PCR assay

The alignment of the three DNA sequences was performed with the program Gene works (Intelligentias, Inc.) by using standard algorithms. Primers were designed to have sequences homologous to those of *A. fumigatus* but not to include the human 18S rRNA gene or the 16S rRNA genes of *Candida* spp. or other pathogenic microorganisms. Therefore, selection of the primer sequences was based on a close check for sequences with matching homologies in current DNA databases (GenBank, release June 1998) with a DNA alignment program (Blast). By using a nested, two-step PCR technique, 4 different primers (Table 1) were tested, and the optimum two pairs (primers AFU5S and AFU5AS and primers AFU7S and AFU7AS) were chosen for all subsequent PCR assays (Figure 1).

**Table 1.** Primer sequence and location.

Primer	DNA sequence (5'-3')	Location	Species specificity
AFU5S	AGG GCC AGC GAG TAC ATC ACC TTG	1436-1459	<i>A. fumigatus</i>
AFU5AS	GG G (AG)GT CGT TGC CAA C(CT)C (CT)CC TGA	1648-1771	<i>A. fumigatus</i>
AFU7S	CGG CCC TTA AAT AGC CCG	1296-1313	<i>A. fumigatus</i>
AFU7AS	GA CCG GGT TTG ACC AAC TTT	1681-1700	<i>A. fumigatus</i>



**Figure 1.** Locations of primer pairs AFU5S-AFU5AS and AFU7S-AFU7AS used in the two-step PCR to detect *Aspergillus* DNA. The primers are derived from the 18S rRNA gene of *Aspergillus* spp. The first PCR step (with AFU7S-AFU7AS) results in amplification of a 405-bp fragment, and the second step with AFU5S-AFU5AS) amplifies an internal fragment of 236 bp.

### Reaction and conditions of (PCR)

Per 25  $\mu$ l PCR mixtures, approximately 50 to 150 ng of total DNA was used as the template. The standard PCR mixture contained 0.5 U of *Taq* DNA polymerase, 6.25 nmol of the deoxynucleoside triphosphates, 10 pmol of primer (first step, primer AFU7S-AFU7AS; second step, primer AFU5S-AFU5AS). In preliminary studies the optimum reaction conditions were established by testing different DNA, primer, enzyme, and deoxynucleoside triphosphate concentrations as well as a range of cycling conditions. PCR was performed in a thermal cycler (Perkin-Elmer Cetus), as follows: for the first PCR, 2 min at 94°C and then 35 cycles of 40 s at 94°C, 1 min at 60°C, and 1 min at 72°C with a terminal step of 5 min at 72°C and then the mixture was held at 4°C; for the second PCR, 2 min at 94°C and then 45 cycles of 40 s at 94°C, 1 min at 60°C, and 1 min at 72°C, with a terminal step of 5 min at 72°C, and then the mixture was held at 4°C. For the second PCR, 10  $\mu$ l of the first-round PCR product was used. The PCR products were separated by 2% agarose gel electrophoresis, stained with ethidium bromide, and visualized with Gel-doc instrument. Control samples included all the constituents in the reaction mixture except genomic DNA. As negative PCR controls, DNA samples of *Phoma*, were used as templates.


### RESULT AND DISCUSSION

The alignment of the 18S rRNA genes of *A. fumigatus*, humans, and *Candida* sp. revealed several regions of significant divergence, which were the basis for selection of the primers. In order to establish a PCR assay specific for several *Aspergillus* species of clinical importance the

18S rRNA gene (rDNA) sequence of *A. fumigatus* (GenBank accession no. AB008401) was aligned with the human 18S rDNA sequence (GenBank accession no. M10098) and the 16S rDNA sequence of *C. albicans* (GenBank accession no. X53497), another ubiquitous microorganism of major clinical importance. Four different primers, comprising 7 sequences upstream and 8 sequences downstream from various divergent regions of the *Aspergillus* species gene, which showed the least homology with the human, or *Candida* genes, were selected for the evaluation and the optimization of the PCR assay.

The two primer pairs that produced PCR products with the highest sensitivity and species specificity were the *A. fumigatus* primers AFU7S and AFU7AS, which amplified a fragment of 405 bp, followed by AFU5S and AFU5AS, which produced an internal fragment of 236 bp (Figures 2 and 3). PCR products could be obtained only from DNA of cultures of *A. fumigatus*, whereas the PCRs with the other *Phoma* test strains were negative. By monitoring the clinical course of this pulmonary tuberculosis patient, in whom a small lung infiltration was detected by chest X ray, we could neither prove nor exclude invasive aspergillosis. PCR has been shown to be a highly sensitive diagnostic tool for the detection of infectious fungi in diverse specimens.

Spradbury et al. (1993) initiated pioneer work in this direction after multicopy gene targets, such as the 28S rRNA gene, had been identified and sequenced. The first PCR assay for the detection of *Aspergillus* DNA in immunocompromised patients described (Melchers et al., 1994) used primers derived from the multicopy 18S rRNA gene, a gene of numerous microorganisms that has now been sequenced. More recent reports described PCR techniques with primers designed to target conserved 18S rRNA sequences common to a variety of fungal pathogens (Einsele et al. 1997; van Burik et al., 1998). However, additional Southern blot analyses with longer probes were also necessary to achieve the detection sensitivity for the identification of the specific fungal species or genera present. Nested PCR assays improved the detection sensitivity. In the first description of such a two-step PCR for the detection of *A. fumigatus* in serum samples of patients, Yamakami et al. (1996) reported that a PCR with two sets of 18S rRNA primers had considerably improved sensitivity compared to that of a PCR assay with a single set of primers. This could be further




**Figure 2.** Determination of the sensitivity of the two-step PCR assay with purified *A. fumigatus* DNA (50 ng). The *Aspergillus* template DNA was clearly detectable by ethidium bromide staining of an agarose gel. Only DNA extracted from *A. fumigatus* (lane 5) was used in a single PCR amplification of the 236-bp fragment with the second primer pair. Negative reagent control amplification without addition of DNA (lane 1 and 2) as well as purified DNA from *Phoma*, DNA resulted in no bands (lane 3, 4). The Lambda DNA EcoR1/Hind III was used as molecular size marker (lane 9). Diluted sample of DNA in (lane 6, 7 and 8) in second step PCR product.

improved by subsequent Southern blot hybridization. These data also highlighted for the first time the use of PCR for the detection of *Aspergillus* DNA in blood samples, which can be obtained by a less noninvasive procedure that is associated with fewer risks than the procedure used to collect BAL specimens, which cannot be performed repeatedly with immunocompromised patients.

We also used a two-step PCR procedure with carefully selected primers to increase the sensitivity and specificity for *A. fumigatus* detection. The multicopy 18S rRNA gene is highly conserved in all species but includes variable regions that are conserved among most *Aspergillus* species. On the basis of comparisons with sequences in a data bank, we chose oligonucleotides from the variable regions that specifically matched only the 18S rRNA gene of *A. fumigatus* to avoid problems arising from co amplification of human or bacterial DNA or contamination with other fungus-derived DNA. Two pairs of nested primers were chosen for use in all subsequent assays. PCR products were obtained only from DNA from *A. fumigatus*.

## Conclusion

A specific two-step PCR assay for the rapid detection of the full range of human pathogenic *Aspergillus* species in sputum samples was established, facilitating improved



**Figure 3.** Determination of the sensitivity of the two-step PCR assay with purified *A. fumigatus* DNA (50 ng). The single derived from 10 fg of *Aspergillus* template DNA was clearly detectable by ethidium bromide staining of agarose gel. As a positive control, only DNA extracted from *A. fumigatus* (10pg) was used in a single PCR amplification of 236-bp fragment with the second primer pair (lane +) A negative control amplification without addition of DNA lane (-) as well as purified DNA from a human cell line (T47D) resulted in no bands. The 123-bp ladder (Gibco BRL ) was used as molecular size marker (lane M ).

early diagnosis and better monitoring of systemic *A. fumigatus* infection in pulmonary tuberculosis patients during therapy in high-risk patients.

## REFERENCES

- Anaissie EJ, Bodey GP, Rinaldi MG (1989). Emerging fungal pathogens. *Eur. J. Clin. Microbiol. Infect. Dis.* 8: 323-330.
- Arning M, Aul C (1994). Prophylaxis against mycosis in neutropenic patients. *Mycoses* 37(2): 70-76.
- Aufavre-Brown A, Cohen J, Holden DW (1992). Use of randomly amplified polymorphic DNA markers to distinguish isolates of *Aspergillus fumigatus*. *J. Clin. Microbiol.* 30: 2991-2993.
- Bart-Delabesse E, Sarfati J, Debeauvais JP, van Leeuwen W, van Belkum A, Bretagne S, Latge JP (2001). Comparison of restriction fragment length polymorphism, microsatellite length polymorphism, and random amplification of polymorphic DNA analyses for fingerprinting *Aspergillus fumigatus* isolates. *J. Clin. Microbiol.* 39: 2683-2686.
- Beck-Sague C, Jarvis WR (1993). Secular trends in the epidemiology of nosocomial fungal infections in the United States, 1980-1990. National Nosocomial Infections Surveillance System. *J. Infect. Dis.* 167: 1247-1251.
- Birch M, Nolard N, Shankland GS, Denning DW (1995). DNA typing of epidemiologically related isolates of *Aspergillus fumigatus*. *Epidemiol. Infect.* 114: 161-168.
- Blum U, Windfuhr M, Buitrago-Tellez C, Sigmund G, Herbst EW, Langer M (1994). Invasive pulmonary aspergillosis. MRI, CT and plain radiographic findings and their contribution for early diagnosis. *Chest.* 106: 1156-1161.
- Bodey GP, Bueltmann B, Duguid W, Gibbs D, Hanak H, Hotchi M, Mall G, Martino P, Meunier F, Milliken S, Naoe S, Okudaira M, Scevola D, van't Wout J (1992). Fungal infections in cancer patients: an international autopsy survey. *Eur. J. Clin. Microbiol. Infect. Dis.* 11: 99-109.
- Bretagne S, Costa JM, Marmorat-Khuong A, Poron F, Cordonnier C, Vidaud M, Fleury-Feith J (2002). Detection of *Aspergillus* species

- DNA in bronchoalveolar lavage samples by competitive PCR. *J. Clin. Microbiol.* 33: 1164-1168.
- Bretagne S, Costa JM, Bart-Delabesse E, Dhedin N, Rieux C, Cordonnier C (1998). Comparison of serum galactomannan antigen detection and competitive polymerase chain reaction for diagnosing invasive aspergillosis. *Clin. Infect. Dis.* 26: 1407-1412.
- Buchheid D, Baust C, Skladny H, Ritter J, Suedhoff T, Baldus M, Seifarth W, Leib-Moesch C, Hehlmann R (2001). Detection of *Aspergillus* species in blood and bronchoalveolar lavage samples from immunocompromised patients by means of 2-step polymerase chain reaction: clinical results. *Clin. Infect. Dis.* 33: 428-435.
- Buchheid D, Baust C, Skladny H, Baldus M, Brauning S, Hehlmann R (2002). Clinical evaluation of a polymerase chain reaction assay to detect *Aspergillus* species in bronchoalveolar lavage samples of neutropenic patients. *Br. J. Hematol.* 116: 803-811.
- Burg JL, Grover M, Pouletty P, Boothroyd JC (1989). Direct and sensitive detection of a pathogenic protozoan, *Toxoplasma gondii*, by polymerase chain reaction. *J. Clin. Microbiol.* 27: 1787-1792.
- Cafferkey MT (1994). Chemoprophylaxis of invasive pulmonary aspergillosis. *J. Antimicrob. Chemother.* 33: 917-924.
- Caillot D, Casanovas O, Bernard A, Couaillier JF, Durand C, Cuisenier B, Solary E, Piard F, Petrella T, Bonnin A, Couillault G, Dumas M, Guy H (1997). Improved management of invasive pulmonary aspergillosis in neutropenic patients using early thoracic computed tomography scan and surgery. *J. Clin. Oncol.* 15: 139-147.
- Chandrasekar PH, Weinmann A, Shearer C (1995). Autopsy-identified infections among bone marrow transplant recipients: a clinicopathologic study of 56 patients. *Bone Marrow Transplant.* 16: 675-681.
- Costa C, Costa JM, Desterke C, Botterel F, Cordonnier C, Bretagne S (2002). Real-time PCR coupled with automated DNA extraction and detection of galactomannan antigen in serum by enzyme-linked immunosorbent assay for diagnosis of invasive aspergillosis. *J. Clin. Microbiol.* 40: 2224-2227.
- Denning DW (1994). Treatment of invasive aspergillosis. *J. Infect.* 28: 25-33.
- Denning DW (1998). Invasive aspergillosis. *Clin. Infect. Dis.* 26: 781-803.
- Einsele H, Hebart H, Roller G, Loeffler J, Rothenhofer I, Muller CA, Bowden RA, van Burik J, Engelhard D, Kanz L, Schumacher U (1997). Detection and identification of fungal pathogens in blood by using molecular probes. *J. Clin. Microbiol.* 35: 1353-1360.
- Einsele H, Steidle H, Vallbracht A, Saal G, Ehninger G, Müller CA (1991). Early occurrence of human cytomegalovirus infection after bone marrow transplantation as demonstrated by the polymerase chain reaction technique. *Blood* 77: 1104-1110.
- Girardin H, Sarfati J, Traore F, Dupouy-Camet J, Derouin F, Latge JP (1994). Molecular epidemiology of nosocomial invasive aspergillosis. *J. Clin. Microbiol.* 32: 684-690.
- Groll AH, Shah PM, Mentzel C, Schneider M, Just-Nuebling G, Huebner K (1996). Trends in the postmortem epidemiology of invasive fungal infections at a university hospital. *J. Infect.* 33: 23-32.
- Guiot HFL, Fibbe WE, van't Wout JWWE (1994). Risk factors for fungal infection in patients with malignant hematologic disorders: implications for empirical therapy and prophylaxis. *Clin. Infect. Dis.* 18: 525-532.
- Hayden RT, Uhl JR, Qian X, Hopkins MK, Aubry MC, Limper AH, Lloyd RV, Cockerill FR (2001). Direct detection of *Legionella* species from bronchoalveolar lavage and open lung biopsy specimens: comparison of *Aspergillus* and *Penicillium* species from respiratory specimens by polymerase of *Aspergillus* infections in a large cohort of patients undergoing bone marrow transplantation. *J. Infect. Dis.* 175: 1459-1466.
- Hearn VM, Pinel C, Blachier S, Ambroise-Thomas P, Grillot R (1995). Specific antibody detection in invasive aspergillosis by analytical isoelectrofocusing and immunoblotting methods. *J. Clin. Microbiol.* 33: 982-986.
- Hebart H, Loeffler J, Reitze H, Engel A, Schumacher U, Klingebiel T, Bader P, Bohme A, Martin H, Bunjes D, Kern WV, Kanz L, Einsele H (2000). Prospective screening by a panfungal polymerase chain reaction assay in patients at risk for fungal infections: implications for the management of febrile neutropenia. *Br. J. Haematol.* 111: 635-640.
- Hendolin PH, Paulin L, Koukila-Kahkola P, Anttila VJ, Malmberg H, Richardson M, Ylikoski J (2000). Panfungal PCR and multiplex liquid hybridization for detection of fungi in tissue specimens. *J. Clin. Microbiol.* 38: 4186-4192.
- Hughes WT, Armstrong D, Bodey GP, Bow EJ, Brown AE, Calandra T, Feld R, Pizzo PA, Rolston KV, Shenep JL, Young LS (2002). Guidelines for the use of antimicrobial agents in neutropenic patients with cancer. *Clin. Infect. Dis.* 34: 730-751.
- Jones ME, Fox AJ, Barnes AJ, Oppenheim BA, Balagopal P, Morgenstern GR, Scarffe JH (1998). PCR-ELISA for the early diagnosis of invasive pulmonary aspergillus infection in neutropenic patients. *J. Clin. Pathol.* 51: 652-656.
- Kami M, Fukui T, Ogawa S, Kazuyama Y, Machida U, Tanaka Y, Kanda Y, Kashima T, Yamazaki Y, Hamaki T, Mori S, Akiyama H, Stevens DA, Kan VL, Judson MA, Morrison VA, Dummer S, Denning D W, Bennett JE, Walsh TJ, Patterson TF, Pankey GA (2000). Practice guidelines for diseases caused by *Aspergillus*. *Infect. Dis. Soc. Am. Clin. Infect. Dis.* 30: 696-709.
- Kami M, Fukui T, Ogawa S, Kazuyama Y, Machida U, Tanaka Y, Kanda Y, Kashima T, Yamazaki Y, Hamaki T, Mori S, Akiyama H, Mutou Y, Sakamaki H, Osumi K, Kimura S, Hirai H (2001). Use of real-time PCR on blood samples for diagnosis of invasive aspergillosis. *Clin. Infect. Dis.* 33: 1504-1512.
- Kappe R, Schulze-Berge A, Sonntag HG (1996). Evaluation of eight antibody tests and one antigen test for the diagnosis of invasive aspergillosis. *Mycoses.* 39: 13-23.
- Kearns AM, Guiver M, James V, King J (2001). Development and evaluation of a real-time quantitative PCR for the detection of human cytomegalovirus. *J. Virol. Methods* 95: 121-131.
- Lass-Flörl C, Aigner J, Gunsilius E, Petzer A, Nachbaur D, Gastl G, Einsele H, Löffler J, Dierich MP, Würzner R (2001). Screening for *Aspergillus* spp. using polymerase chain reaction of whole blood samples from patients with hematological malignancies. *Br. J. Haematol.* 113: 180-184.
- Lewin SR, Vesanen M, Kostrikis L, Hurley A, Duran M, Zhang L, Ho DD, Markowitz M (1999). Use of real-time PCR and molecular beacons to detect virus replication in human immunodeficiency virus type 1-infected individuals on prolonged effective antiretroviral therapy. *J. Virol.* 73: 6099-6103.
- Lin S, Schranz J, Teutsch S (2001). Aspergillosis case-fatality rate: systematic review of the literature. *Clin. Infect. Dis.* 32: 358-366.
- Loeffler J, Hebart H, Brauchle U, Schumacher U, Einsele H (2000a). Comparison between plasma and whole blood specimens for detection of *Aspergillus* DNA by PCR. *J. Clin. Microbiol.* 38: 3830-3833.
- Loeffler J, Henke N, Hebart H, Schmidt D, Hagemeyer L, Schumacher U, Einsele H (2000). Quantification of fungal DNA by using fluorescence resonance energy transfer and the Light Cycler system. *J. Clin. Microbiol.* 38: 586-590.
- Loeffler J, Hebart H, Sepe S, Schumacher U, Klingebiel T, Einsele H (1998). Detection of PCR-amplified fungal DNA by using a PCR-ELISA system. *Med. Mycol.* 36: 275-279.
- Logan PM, Primack SL, Miller RR, Muller NL (1994). Invasive aspergillosis of the airways: radiographic, CT and pathologic findings. *Radiology.* 193: 383-388.
- Makimura K, Murayama SY, Yamaguchi H (1994). Detection of a wide range of medically important fungi by the polymerase chain reaction. *J. Med. Microbiol.* 40: 358-364.
- Melchers WJ, Verweij PE, van den Hurk P, van Belkum A, De Pauw BE, Hoogkamp-Korstanje JA, Meis JF (1994). General primer-mediated PCR for detection of *Aspergillus* species. *J. Clin. Microbiol.* 32: 1710-1717.
- Obayashi T, Yoshida M, Mori T, Goto H, Yasuoka, Iwasaki A, HTeshima H, Kohno S, Horiuchi A, Ito A (1994). Plasma (1-3)-beta-D-glucan measurement in diagnosis of invasive deep mycosis and fungal febrile episodes. *Lancet* 345: 17-20.
- Palladino S, Kay I, Fonte R, Flexman J (2001). Use of real-time PCR and the LightCycler system for the rapid detection of *Pneumocystis carinii* in respiratory specimens. *Diagn. Microbiol. Infect. Dis.* 39: 233-236.
- Pfaffenbach B, Donhuijsen K, Pahnke J, Bug R, Adamek R, Wegener

- JM, Ricken D (1994). Systemische Pilzinfektionen bei hämatologischen Neoplasien. Eine Autopsiestudie and 1053 Patienten. Med. Klin 89: 299-304.
- Skladny H, Buchheidt D, Baust C, Krieg-Schneider F, Seifarth W, Leib-Mosch C, Hehlmann R (1999). Specific detection of *Aspergillus* species in blood and bronchoalveolar lavage samples of immunocompromised patients by two-step PCR. J. Clin. Microbiol. 37: 3865-3871.
- Spreadbury C, Holden D, Aufavre-Brown A, Bainbridge B, Cohen J (1993). Detection of *Aspergillus fumigatus* by polymerase chain reaction. J. Clin. Microbiol. 31: 615-621.
- Stynen D, Goris A, Sarfati J, Latge JP (1995). A new sensitive sandwich enzyme-linked immunosorbent assay to detect galactofuran in patients with invasive aspergillosis. J. Clin. Microbiol. 33: 497-500.
- Swanink CMA, Meis JFGM, Rijs AJMM, Donnelly JP, Verweij PE (1997). Specificity of a sandwich enzyme linked immunosorbent assay for detecting *Aspergillus* galactomannan. J. Clin. Microbiol. 35: 257-260.
- van Burik JA, Myerson D, Schreckhise RW, Bowden RA (1998). Panfungal PCR assay for detection of fungal infection in human blood specimens. J. Clin. Microbiol. 36: 1169-1175.
- Verweij PE, Rijs AJ, De Pauw BE, Horrevorts AM, Hoogkamp-Korstanje JA, Meis JF (1995). Clinical evaluation and reproducibility of the Pastorex *Aspergillus* antigen latex test for diagnosing invasive aspergillosis. J. Clin. Pathol. 48:474-476.
- Wald A, Leisenring W, Van Burik JA, Bowden RA (1997). Successful Nonmyeloablative Allogeneic Hematopoietic Stem Cell Transplant in an Acute Leukemia Patient With Chemotherapy-Induced Marrow Aplasia and Progressive Pulmonary *Aspergillosis* Blood, Epidemiology 94: 3273-3276.
- Walsh TJ, Francesconi A, Kasai M, Chanock SJ (1995a). PCR and single-strand conformational polymorphism for recognition of medically important opportunistic fungi. J. Clin. Microbiol. 33: 3216-3220.
- Walsh TJ, Garrett K, Feuerstein E, Girton M, Allende M, Bacher J, Francesconi A, Schaufele R, Pizzo PA (1995b). Therapeutic monitoring of experimental invasive pulmonary aspergillosis by ultrafast computerized tomography: a novel, noninvasive method for measuring responses to antifungal therapy. Antimicrob. Agents Chemother. 39: 1065-1069.
- Walsh TJ, Lee J, Lecciones J, Rubin M, Butler K, Francis P, Weinberger M, Roilides E, Marshall D, Gress J, Pizzo PA (1991). Empiric therapy with amphotericin B in febrile granulocytopenic patients. Rev. Infect. Dis. 13: 496-503.
- Yamakami Y, Hashimoto A, Tokimatsu I, Nasu M (1996). PCR detection of DNA specific for *Aspergillus* species in serum of patients with invasive aspergillosis. J. Clin. Microbiol. 34: 2464-2468.