

Full Length Research Paper

# HIV-Malaria coinfection in the Department of Paediatrics of Centre Hospitalier Universitaire Sourou SANOU ( CHUSS )

Ouedraogo SM<sup>1,2</sup>, Sangaré I<sup>1,2</sup>, Sourabié Y<sup>1,2</sup>, Zongo R<sup>1</sup>, Zida A<sup>3,4</sup>, Ouedraogo AS<sup>1,2</sup>, Fofana S<sup>1,2</sup>, Guiguemdé RT<sup>2</sup>

<sup>1</sup>Sourou Sanou University Hospital, Mail box 676 Bobo-Dioulasso 01 Burkina Faso

<sup>2</sup>Higher institute of Health sciences, Polytechnic University of Bobo Dioulasso, Burkina Faso.

<sup>3</sup>University department of Health sciences, University of Ouagadougou, Burkina Faso.

<sup>4</sup>Yalgado Ouedraogo University Hospital, Ouagadougou, Burkina Faso.

Accepted 28 April, 2015

**Introduction:** The interaction of these two pathologies raises capital issues as well as therapeutic. This study aims at a better understanding of the clinical, biological and therapeutic profile of HIV-malaria coinfecting children.

**Methodology:** It is a prospective and retrospective cross-sectional study. Were involved patients infected with HIV followed in the department whose clinical and laboratory diagnosis of malaria had been made and / or patients with whom the diagnosis of coinfection was made through a medical consultation.

**Results:** The prevalence of coinfection was 3.09%. The clinical signs of severity were dominated by impaired consciousness and seizures. As for biological signs, anemia was found with 65% of patients and thrombocytopenia with 9.3% of cases. Immunocompetent patients accounted for 70.7% of coinfecting patients. Pharmacological therapeutic groups which were administered were hyperleucocytoses, they were found with 19.6% of coinfecting patients mainly antimalarials (100% of cases), analgesics-antipyretics (79.6% of cases) and ARV(48.1% of cases).

**Conclusion:** Immunosuppression caused by HIV infection did not appear to be associated with the frequency of occurrence of malaria. Chemo malaria prophylaxis with HIV positive children is not necessary.

**Key words:** Malaria, HIV, children, Bobo-Dioulasso, Burkina Faso.

## INTRODUCTION

HIV and Malaria infection are nowadays two important public health issues in sub-Saharan Africa. They are the leading causes of morbidity and mortality. Indeed this coinfection (HIV and malaria) bring about 4 million deaths every year in developing countries. Each year a minimum of 50 million pregnancies are recorded with women living in malaria-endemic areas, with about half of them in sub-Saharan Africa. In this part of Africa, it is estimated that one million pregnancies each year occur with women suffering from an HIV- malaria coinfection (Cuadros et al., 2011).

Coinfection is much found with children in this part of the

world, since vertical transmission is the main route of HIV transmission to children. In fact, according to statistics, a child is subject to at least one fit of malaria during each transmission season. Malarial transmission is prevalent in the endemic-epidemic mode with an increase in the rainy season (from July to October). With regards to HIV infection, there were 10,000 children infected with HIV, living in Burkina Faso in 2008. Bobo-Dioulasso is among the cities in Burkina Faso where the prevalence of HIV infection is high. In 2007 it was estimated at 3%, while the national prevalence was 2%. The profound impact of these infections on the immune system is generating interest in the scientific community (Renia et al., 2006). Indeed, previous studies reported that HIV infection would be responsible for an increased incidence of malaria attacks and an increase in parasitaemia (Blondel et al., 2007; Grimwade et al., 2004; Diallo et al., 2004; Verhoeff et al. 1999). However, this finding is not shared

by all; in fact, some authors have been reporting low malaria prevalence (11.4%) with children infected with HIV compared to HIV-negative children (27.6%) (Villamor et al. 2003). HIV- malaria coinfection may also raise diagnostic issues. Indeed, for some authors high parasitaemia would be associated with a high prevalence of "false positive" of immune-enzyme tests, especially with young patients (Gasasira et al., 2006; Gosh et al., 2001). Furthermore, therapeutic issues were raised by some authors (Shah et al., 2006; Kanya et al., 2006).

Thus, the combination of therapies based on artemisinin (CTA) would have potential interactions with Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs) and protease inhibitors (PIs) that are metabolized by CYP450 (Alemu et al. 2013). Also, the co-administration of sulfadoxine-pyrimethamine combination (SP) with cotrimoxazole or nevirapine increases the risk of liver and skin side effects, and with zidovudine, the risk of bone marrow toxicity.

Evolutionary interactions between these two diseases have been reported by other authors. In fact, during the acute phase of malaria the viral load of HIV would raise, increasing the risk of transmission, resulting in a decline in CD4 rate (Kublin et al., 2005; Slutsker et al., 2007).

This study analyzed the clinical, biological and therapeutic profile of HIV-malaria coinfection with children followed at Hospital du Jour of Centre Hospitalier Universitaire Sourou SANOU (CH SS), of Bobo-Dioulasso in order to allow a better understanding of the interaction between these two major diseases in Burkina Faso.

## METHODOLOGY

The study was carried out in the Department of Pediatrics of CH SS, Bobo-Dioulasso in Western Burkina Faso. Two types of studies were carried out from July to December 2013:

- 1- A Prospective cross-sectional study about children whose HIV status was unknown at admittance;
- 2- A retrospective cross-sectional study about children infected with HIV and followed in the department

Population 1 consisted of HIV positive patients, followed in the department and whose clinical and laboratory diagnosis of malaria was made.

Population 2 consisted of patients who consulted the pediatric department for a suggestive clinical picture of malaria confirmed by parasitological examination; those with positive HIV serology were involved. The involvement was made after obtaining the informed and free consent of parents or legal guardians.

For population 1 data were collected from the physical and electronic records of patients. And for population 2, the clinical diagnosis was provided by a clinician, and laboratory confirmation of malaria was made from thick blood smears after coloring with Giemsa 10% and viewed by optical microscope. Positive cases in Search for

Plasmodium (SP) were submitted to a rapid HIV screening (Determine®). A confirmation and typifying test was done when the first test was positive. SD Bioline® was used for this purpose. Sociodemographic, clinical, biological, pharmacological therapy data were collected using a questionnaire.

Newborns were aged from 0 to 29 days and infants aged from 1 to 30 months, the toddlers over 30 months to 5 years, as for late toddlers and teenagers their proportion of respective age was from over 5 years to 10 years and from over 10 years to 18 years. We considered severe anemia an hemoglobin rate (Hb) less than 7 g / L; moderate anemia an Hb rate between 7 and 9 g / L and a light anemia, an Hb rate, ranging from beyond 9 g and less than or equal to 11g / L. As for CD4 rate, we have taken into account the WHO recommendations of putting on treatment (2010), to classify our patients as follows:

- Children 0-59 months [0-750 [cell /  $\mu$ l = Immunocompromised, 750 and more = immunocompetents.
- Children aged 60 months and over: [0-350 [cell /  $\mu$ l = Immunocompromised, 350 and more = immunocompetents.

The statistical test used was the Chi square test at 5% significance threshold. For low numbers and in cases where one of the expected values in a cell was less than 5, Yates' correction was applied.

## RESULTS

### Epidemiological aspects of HIV- Malaria coinfection

In population 1: Involved, were children infected with HIV and followed in the department. The number of visits during the study period was 3313 including 102 cases of malaria, i.e. a coinfection prevalence of 3.07%.

In population 2: Involved were Children who were diagnosed with malaria at admittance whose HIV status was determined when hospitalized. Out of 181 cases, there were 6 patients infected with HIV and 4 patients put at risk (patients under 18 months who were tested positive to the rapid test: Determine®) i.e, a prevalence of 3.3% in this population.

The overall prevalence of coinfection was estimated at 3.09%.

### Distribution of coinfection per age

The prevalence was higher in the group of teenagers (43.5%), Figure 1 shows the distribution of coinfection per age

### Monthly dynamics of coinfection

Malaria and HIV coinfection was more observed during malaria transmission season (July-October). It was




Figure 1. Number of cases of co infection per child group age.




Figure 2. Distribution of cases of co infection with the stage of HIV infection and evolution.

12.9% (14 cases) in July, 20.4% (20 cases) in August, 20.4% (22 cases) in September and 27.8% in October i.e., 30 cases. Respective rates of 12% (13 cases) and 8% (9cas) were observed in November and December.

**Clinical aspects of coinfection**

The most common clinical signs in the 2 populations were fever, headache and vomiting. Signs of seriousness were dominated in population 1 by consciousness

disorders (3.9%), hemoglobinuria (0.98%), convulsion (0.98%) and jaundice (0.98%), whereas in population 2 we noted dehydration (50%), disorders of consciousness (33.3%), coma (16.7%) and seizures (33.3%).

**Distribution of patients in accordance with the type of malaria**

Severe cases of malaria were 14% versus 86% for simple malaria. All patients diagnosed HIV positive at




Figure 3. Distribution of cases of co infection based on the type of anemia.




Figure 4. reports the distribution of different combinations of ARTs used.

admittance were suffering from severe malaria, while with children infected with HIV and followed at the Pediatrics Department, severe malaria concerned only 8.8% of cases of coinfection.

**Classification of patients in accordance with WHO clinical stage**

Figure 2 shows the frequency of different clinical stages of HIV infection in accordance with WHO classification. Patients in stage 1 represented the majority of cases

(60.6%). There was a significant difference between clinical stage and evolution. The risk of death was higher with patients at advanced stages.

**Biological aspects of coinfection**

In both populations as shown in Figure 3, there was a higher incidence of mild anemia. In population 1, there was 66.7% of mild anemia; 22.8% of moderate anemia and 10.5% of severe anemia. In Population 2, severe anemia was found in 66.6% of

**Table 1.** Frequency of therapeutic groups prescription.

<b>pharmacological Groups</b>	<b>Frequency</b>	<b>Percentages</b>
<i>Antimalarials</i>	108	100%
<i>Antalgesics-Antipyretics</i>	86	79.6%
<i>ART</i>	52	48.1%
<i>Antibiotics and others</i>	30	27.8%
<i>Cotrimoxazole</i>	23	21.3%
<i>Antiemetics</i>	12	11.1%
<i>Antitussives</i>	12	11.1%
<i>Antihistamines</i>	5	9.3%
<i>Blood and derivatives</i>	4	3.7%
<i>Iron</i>	3	2.8%
<i>Oral solutes</i>	3	2.8%
<i>Antispasmodics</i>	2	1.9%
<i>Parasites killers</i>	2	1.9%
<i>Anticonvulsants/sedatives</i>	1	0.9%

patients with an equal distribution of the other two types of anemia i.e., 16.7%.

Leukocytosis was found with 19 patients versus 7 cases of leukopenia. With 71 patients no abnormalities of the leukocyte rate was observed.

In Population 2, 50% of patients had leukocytosis versus 17.6% in population 1. 16 patients were suffering from thrombocytosis and 9 from thrombocytopenia.

The majority of patients showed no platelet defects (74.2%).

### **Immunological data**

#### **Distribution of patients in accordance with their immunological profile**

The frequency of HIV-1 infection was 97.2%. We reported a rate of 29.3% of immunocompromised versus 70.7% of immunocompetent. Before 60 months there was a neat predominance of immunosuppression (71.4%), conversely after 60 months, immunocompetence predominated (75%).

#### **Distribution of patients in accordance with the CD4 rate and type of malaria**

Children under 60 months suffered all from uncomplicated malaria. With the 60 months old and more, uncomplicated malaria had an almost equal distribution between immunocompetent and immunocompromised with respective rates of 90% and 88%. However, severe malaria was found with at a higher

frequency with immunocompromised (11.8%) versus 9.8% with immunocompetent.

### **Pharmacotherapeutic support**

Antimalarials were prescribed to all patients at respective rates of 86.1% (93/108) and 13.9% (15/108) for amodiaquine / artesunate and quinine, and 48.1% of patients were on antiretroviral therapy (ART).

Table 1 shows the distribution of pharmacotherapeutic groups used in coinfection.

As for ARV treatment, the combination of d4T-3TC-NVP represented the group of ARVs used mostly. Figure 4 reports the distribution of the various combinations of used ART

### **Discussion**

#### **Prevalence of coinfection**

The overall prevalence was 3.09%. This rate is lower than those reported by Villamor, Tanzania (11.4%) and Whitworth, Uganda (11.8%) (Villamor et al., 2003; Whitworth et al., 2000). However, these two studies estimated the existence of parasitaemia or not with patients infected with HIV and not malaria. Thus the prevalence of malaria in their respective studies could be lower because asymptomatic carriage of the parasite in stable endemic countries is common.

The low prevalence of malaria in children infected with HIV is also explained by the compliance with a reinforcement of medical care with these patients, and

sensitization of parents and guardians to early medical consultation at the slightest symptom of clinical suspicion of malaria. Kawo shares our results in his study by relating malaria prevalence 4 times higher with HIV-negative children compared to children infected with HIV (Kawo et al., 2000).

Several hypotheses were put forward to explain this finding including:

- Co-trimoxazole prophylaxis, which would significantly reduce the risk of occurrence of malaria in children infected with HIV. In Mali, Thera reported a reduction in the incidence of malaria cases with children under chemo prophylaxis not infected with HIV. Such a finding is shared by several other authors who reported in their work a reduction of the incidence of malaria attacks and mortality among HIV-infected patients on cotrimoxazole (Shah et al., 2006; Slutsker et al., 2007; Mermin et al., 2006; Saracino et al., 2012; Lockwood et al., 2004);
- Oxidative stress created by HIV infection in the erythrocytes that would be unfavorable to the development of Plasmodium (Chaisavaneeyakorn et al., 2002);
- Antiretroviral therapy would promote the control of malaria infections, firstly, by restoring the antimalarial immune response and on the other, by the effect of some molecules that could have an antimalarial effect *in vitro*, it is the case of protease inhibitors such as ritonavir and saquinavir which inhibit *in vitro* the parasite growth at clinically effective concentrations (Davenport et al. 2012; Skinner-Adams et al. 2004). So, ARV treatment by non-nucleoside reverse transcriptase inhibitors would be associated with a decrease in the incidence of malarial attacks without ruling out the involvement of the effects of immune restoration (Davenport et al; 2010). Thus Mermin brought through his studies an impact of ARV treatment with coinfecting patients. Indeed, the annual incidence of malaria without intervention was 50.8 episodes per 100 people per year, which was reduced to 9 malaria episodes per 100 people per year with chemoprophylaxis and cotrimoxazole only and 3.5 episodes per 100 people per year with cotrimoxazole and antiretroviral treatment (Mermin et al., 2006).
- The reduction of HIV replication in secondary human macrophages resulting from the presence of Plasmodium haemozoin in endothelial cells of the malaria cured patients or with chronic infection. *In vitro*, indeed, this reduction is associated with a restriction on the integration of the viral genome into the chromosome of the target host cells (Davenport et al; 2010).

#### Distribution of coinfection in accordance with age

In **population 1**, the prevalence of co-infection was low with those under 24 months. The administration of cotrimoxazole with prophylactic perspective being systematic with HIV-exposed children up to the age of 18 months before showing their non-contamination, could

justify this finding.

However in **Population 2**, children from 0-5 years old accounted for 83% of malaria-stricken population. The high incidence of malaria in this age group is well known in areas of endemic malaria prophylactic and extensive sensitization campaigns are directed towards this target group in Burkina (Oura, 2009).

#### Temporal distribution of co-infection

The monthly dynamics of malaria transmission in Bobo-Dioulasso may justify the increasing prevalence of malaria from July to October. Indeed in western Burkina Faso, malaria is rampant in an endemic- epidemic mode with a high prevalence during the rainy season (July to September) and a peak in early dry season (October) (Baudon et al., 1988).

#### Clinical profile

##### Clinical signs

Clinically, the three main reasons for consultation in **population 1** were pyrexia (97%), headache (61.8%) and vomiting (37.3%). Within this population signs of severity were pallor (3.9%), disorders of consciousness (3.9%), repeated convulsions (0.98%), hemoglobinuria (0.98%) and jaundice (0.98%). In **Population 2**, the signs of severity differed from those observed in population 1 with higher proportions of dehydration (50%), repeated convulsions (50%), disorders of consciousness (33.3%) of pallor (33.3%), coma (16.7%). Regular monitoring of patients infected with HIV and the effectiveness of management protocols for these children explain the low levels of seriousness signs in population 1. Although these signs are often found in the symptomatology of severe malaria regardless of any HIV infection (Maitland et al., 2004), this difference in distribution of seriousness signs in children infected with HIV and followed, and children whose HIV status was discovered at the moment of suspected signs of malaria should be elucidated (Greenberg et al., 1988; Nguyen-Dinh et al., 1987).

##### WHO clinical stage

The majority of patients were in stage 1 of HIV infection (56%) and death was significantly correlated with WHO clinical stage which was high at the advanced stage of HIV infection. However, the accountability of malaria as a cause of death at this late stage of HIV infection when opportunistic infections are predominant remains difficult to confirm.

##### Biological profile

##### Hematology data

In population 1, mild anemia, however, was predominant

in population 2; severe anemia was reported in more than half of this population (66.7%).

The strong correlation severe anemia-HIV infection-severe malaria (whatever the parasite density) in a high transmission area of malaria is also reported by several other authors such as Oura and Otieno and could explain the high prevalence of severe anemia in the population where the severe form of malaria prevailed (Oura, 2009 ; Otieno et al. 2002).

In both populations, leukocytosis was the most frequent quantitative abnormality of leukocytes. The majority of patients exhibited no quantitative platelet defects (74.2%).

### Immunological data

HIV Serotype 1 was predominant and found in 97.2% in our study. The prevalence of this serotype in Africa is known and reported by several authors (Whitworth et al., 2000; Kawo et al., 2000). As it is clear from our series, a high prevalence of immunocompetent individuals (70.7%).

Yet many authors through their studies have reported a drop in CD4 rates during a malaria attack, Mermin and Whitworth, Uganda (Whitworth et al., 2000; Mermin et al., 2006). The unavailability of CD4 rate before malaria with some of our patients (population 2), does not permit the establishment of any link of changes in CD4 levels before, during and after an attack of malaria. But no relationship was observed between the clinical form of malaria and the immunological profile of patients.

### Therapeutic profile

The most prescribed antimalarials were the combination amodiaquine - artesunate with a rate of 86.1%. This is explained by the predominance of the simple form of malaria, the diagnosis was made with 86% of our patients, indeed, in accordance with the national guidelines in Burkina Faso, which is part of the strategic plan to fight against malaria (2011-2015), the treatment of uncomplicated malaria uses the combination therapy amodiaquine - artesunate or artemether-lumefantrine by oral route; injectable forms of artesunate, artemether and quinine are reserved for severe forms which concerned only 14% our patients (Oura, 2009).

Co infection does not require changing the therapeutic protocol compared to the treatment of uncomplicated malaria and severe malaria, which is shared by (Brentlinger et al. 2007), although for some authors a delay was found in the clearance of parasite with patients infected with HIV and treated by means of artemisinin, suggesting a metabolic disorder of antimalarials according to the immunological profile of the host (Birku et al., 2002).

### CONCLUSION

This study showed a low prevalence of HIV-malaria co-

infection with children. But malaria can be a testing ground for HIV infection. Coinfection involved more teenagers than infants with a strong higher frequency during malaria transmission season. Leukocytosis is the most common laboratory abnormality during this coinfection. No relationship was observed between the clinical form of malaria and the immunological profile of patients.

The treatment of malaria during coinfection followed national guidelines for the fight against malaria. Signs of severe malaria were less observed in children infected with HIV and followed in pediatrics.

Therefore, devolution of health structures monitoring children infected with HIV, involving Health Centers and Social Promotion (CSPS), combined with a strengthening of medical monitoring of therapeutic and technical capacities of children infected with HIV- malaria is essential.

### CONFLICTS OF INTERESTS

**Authors report no conflict of interest opposing them**

### REFERENCES

- Alemu A, Shiferaw Y, Addis Z, Mathewos B, Birhan W (2013). Effect of malaria on HIV/AIDS transmission and progression. *Parasit Vectors*; 6:18.
- Baudon D, Galaup B, Ouedraogo L, Gazin P (1988). Une étude de la morbidité palustre en milieu hospitalier au Burkina Faso (Afrique de l'Ouest). *Med Trop.*, 48:9-13.
- Birku T, Mekonnen E, Bjorkman A, Wolday D (2002). delayed clearance of Plasmodium falciparum in patients with HIV virus coinfection treated with artemisinin. *Ethiop MedJ.*, 40:17-26.
- Blondel G, Barry M, Porte L (2007). Impact de l'infection à VIH sur l'infection palustre chez l'adulte. *Med et Mal In.*, 37:629-636.
- Brentlinger PE, Behrens CB, Kublin JG (2007). Challenges in the prevention, diagnosis, and treatment of malaria in human immunodeficiency virus infected adults in sub-Saharan Africa. *Arch Intern Med.*, 167:1827-1836.
- Chaisavaneeyakorn S, Moore JM, Otieno J, (2002). Immunity to placental malaria. III. Impairment of interleukin(IL)-12, not IL-18, and interferon-inducible protein-10 responses in the placental intervillous blood of human immunodeficiency virus/malaria-coinfected women. *J Infect Dis.*, 185:127-131.
- Cuadros DF, Branscum AJ, Crowley PH (2011). HIV-malaria co-infection: effects of malaria on the prevalence of HIV in East Sub-Saharan Africa. *Int J Epidemiol*; 40:931-939.
- Davenport GC, Hittner JB, Were T, Ong'echa JM, Perkins DJ (2012). Relationship between inflammatory mediator patterns and anemia in HIV-1 positive and exposed children with Plasmodium falciparum malaria. *Am J Hem-*

*atol.*,87:652-658.

Davenport GC, Ouma C, Hittner JB (2010). Hematological predictors of increased severe anemia in Kenyan children coinfecting with Plasmodium falciparum and HIV-1. *Am J Hematol.*, 85:227-233

Diallo AH, Ki-Zerbo G, Sawadogo AB, Guiguemde TR (2004). Paludisme grave et infection à VIH chez l'adulte à Bobo-Dioulasso, Burkina Faso. *Med Trop.*, 64:345-350.

Gasasira A, Dorsey G, Kanya M, Havlir D, Kiggundu M, Rosenthal P et al. (2006). False-positive results of enzyme immunoassays for human immunodeficiency virus in patients with uncomplicated malaria. *J Clin Microbiol.*, 44:3021-3024.

Gosh K, Javeri K, Mohanty D, Parmar B, Surati R, Joshi S (2001). False positive serological tests in acute malaria. *Br J Biomed Sci.*, 58:20-23.

Greenberg A, Nguyen-Dinh P, Mann J, Kabote N, Colebunders R, Francis H (1988). The association between malaria, blood transfusions, and HIV seropositivity in a pediatric population in Kinshasa, Zaire. *JAMA.*, 259:10-11.

Grimwade K, French N, Mbatha D (2004). HIV infection as a cofactor for severe falciparum malaria in adults living in a region of unstable malaria transmission in south africa. *AIDS.*, 18:547-554.

Kanya M, Gasasira A, Yeka A (2006). Effect of HIV-1 infection on antimalarial treatment outcomes in Uganda : a population-based study. *J Infect Dis.*, 193:9-15.

Kawo G, Karlson K, Lyamuya E (2000) Prevalence of HIV type 1 infection associated clinical features and mortality among hospitalized children in Dar es Salaam, Tanzania. *Scand J Infect Dis.*, 32:357-363.

Kublin JG, Patnaik P, Jere C (2005). Effect of Plasmodium falciparum malaria on concentration on HIV-1 RNA in the blood of adults in rural Malawi : a prospective cohort study. *Lancet*, 365:233-240.

Lockwood D, Armstrong M, Grant A (2004). Integrating evidence based medicine into routine clinical practice: seven years' experience at the Hospital for Tropical Diseases, London. *BMJ.*, 329:1020-1023.

Maitland K, Marsh K (2004). Pathophysiology of severe malaria in children. *Acta Trop.*, 90:131-140.

Mermin J, Ekwaru J, Liechty C, Were W, Downing R, Ransom R (2006). Effect of cotrimoxazole prophylaxis, antiretroviral therapy, and insecticide-treated bed nets on the frequency of malaria in HIV-1 infected adults in Uganda : a prospective cohort study. *Lancet*, 367:1256-1261.

Mermin J, Lule J, Ekwaru J (2006). Association between malaria and CD4 cell count decline among persons with HIV. *J Acquir Immune Defic Syndr.*, 41:129-130.

Nguyen-Dinh P, Greenberg A, Mann J (1987). Absence of association between P. falciparum malaria and HIV infection in children in Kinshasha (Zaire). *Bull OMS.*, 65:607-613.

Otieno R, Ouma C, Ong'echa JM (2006). Increased severe anemia in HIV-1 exposed and HIV-1 positive infants and children during acute malaria. *AIDS.*, 20:275-280.

Oura M (2009). Problématique du diagnostic de gravité du paludisme dans un service de référence. *Université de Ouagadougou*:180p.

Renia L, Potter S (2006). Coinfection of malaria with HIV : an immunological perspective. *Parasite immunol.*, 28:589-595.

Saracino A, Nacarapa EA, da Costa Massinga EA, (2012) Prevalence and clinical features of HIV and malaria co-infection in hospitalized adults in Beira, Mozambique. *Malar J.*, 11:241.

Shah S, Smith E, Obonyo C, Kain KC, Bloland P, Slutsker L et al. (2006) HIV immunodepression and antimalarial efficacy : sulfadoxine-pyrimethamine for the treatment of uncomplicated malaria in HIV-infected adults in Siaya, Kenya. *J Infect Dis.*, 194:1519-1528.

Skinner-Adams TS, McCarthy JS, Gardiner DL, Hilton PM, Andrews KT (2004). Antiretrovirals as antimalarial agents. *J Infect Dis.*, 190:1998-2000.

Slutsker L, Marston BJ (2007). HIV and malaria: interactions and implications. *Curr Opin Infect Dis.*, 20:3-10.

Thera M, Sehdev P, Coulibaly D (2005). Impact of trimethoprim-sulfamethoxazole prophylaxis on falciparum malaria infection and disease. *J Infect Dis.*, 192:1823-1829.

Verhoeff F, Brabin B, Hart C, Chimsuku L, Kazembe P, Broadhead R (1999). Increased prevalence of malaria in HIV-infected pregnant women and its implications for malaria control. *Trop Med Int Health.*, 4:5-12.

Villamor E, Fataki M, Mbise R, Fawzi W (2003). Malaria parasitaemia in relation to HIV status and Vitamin A supplementation pre-school children. *Trans R Soc Trop Med Hyg.*, 8:1051-1061.

Whitworth J, Morgan D, Quigley M (2000) Effect of HIV-1 and increasing immunosuppression on malaria parasitaemia and clinical episode in rural Uganda : a cohort study. *Lancet*, 356:1051-1056.