

Full Length Research Paper

# Assessment of the genetic diversity in five generations of a commercial broodstock line of *Litopenaeus vannamei* shrimp

Patrícia Domingues de Freitas and Pedro Manoel Galetti Jr.

Departamento de Genética e Evolução, Universidade Federal de São Carlos, Caixa Postal 676, 13565-905 - São Carlos, SP, Brazil (correspondence to P.D.F.)

Accepted 12 August, 2019

Genetic variation among and within five generations of an inbred commercial captive line of *Litopenaeus vannamei* and genetic distance among them were evaluated by random amplified polymorphic DNA (RAPD), using descriptive and genetic similarity analyses for dominant markers at single- and multi-population level. One hundred individuals were analyzed and 56 polymorphic loci were identified with a set of six primers screened in a panel of 38 decamer primers tested. The values obtained for F<sub>5</sub>, F<sub>6</sub>, F<sub>7</sub>, F<sub>8</sub> and F<sub>9</sub> generations revealed a progressive increasing of genetic similarity throughout the five stocks, confirmed by Nei genetic diversity analysis, which values decreased from 0.27 to 0.22 at F<sub>5</sub> and F<sub>9</sub>, respectively. Significant allele frequency differences were observed at most analyzed loci. Strong correlation ( $R^2 = 0.9845$ ) was observed between genetic similarity and generation time, and genetic similarity could reach closer to 100% at 18-19 generation of this line.

**Key words:** *Litopenaeus vannamei*, genetic diversity, genetic drift, inbreeding, RAPD.

## INTRODUCTION

Loss of genetic variation in small populations can be a consequence of genetic drift and inbreeding, commonly observed within captive and/or threatened stocks (Sbordoni et al., 1986). The low levels of genetic variability may reduce the mean fitness of a population, affecting its viability, particularly, if parasites or competitors are present (Vrijenhoek, 1994). It is well known that the genetic variability declining in natural populations, associated to inbreeding and genetic drift, increases the probability of extinction of small populations, determining, in some cases, a process called extinction vortex in nature (Gilpin and Soulé, 1986).

Within captive populations, these effects are far well known (Bradford et al., 1958), including in aquaculture-

based larger effective populations (Tave, 1993). It is important for a successful aquaculture to prevent such genetic variation losses.

Genetic variability reduction has previously been reported in reared shrimp. In *Marsupenaeus japonicus*, both inbreeding and genetic drift were implicated in a continuous reduction in the genetic diversity levels throughout five captive generations (Sbordoni et al., 1986).

It is already recognized that the current Brazilian shrimp farming activity involves an excellent management technology, with high productivity around 6.5 ton/ha/year (Rocha et al., 2004). Several advances were made after the introduction of the Pacific white shrimp, *Litopenaeus vannamei*, currently corresponding to 90% of all farmed shrimp in Brazil. The complete control of the reproductive cycle and post-larvae production have consolidated the Brazilian broodstocks, making it unnecessary for the importation of post-larvae and adult prawns, and avoiding the introduction of exotic pathogens (Batalha et al., 2002). Apart from the remarkable development of shrimp culture in

\*Corresponding authors E-mail: [patdf@iris.ufscar.br](mailto:patdf@iris.ufscar.br);  
Tel: 55 16 2608309; Fax: 55 16 260 8306.

Brazil, selective breeding programs have poorly been conducted (Galetti and Freitas, 1999). Most of hatchery laboratories develop management programs without any monitoring of the levels of genetic variability of their broodstocks.

In the present study, five generations of a closed line of the white-shrimp *L. vannamei* were analyzed by RAPD markers to assess the genetic variation within and between the cultured generations, as well as to infer the more likely causes leading to genetic variability reduction in these animals.

## MATERIAL AND METHODS

### Sample collection

A commercial line of *L. vannamei* owned by Valença da Bahia Maricultura shrimp hatchery and farm, located on the Northeastern region of Brazil in the municipality of Valença (Bahia state), was studied for five generations. The founder individuals of this line were obtained in 1995 by importation of native individuals mainly from Panama (80%), and native and farmed individuals from other countries including Ecuador, Venezuela, Costa Rica, and Mexico (20%). When adapted to captivity conditions, males and females were selected and, after the first spawning on captive conditions, the generation F<sub>1</sub> was obtained. At each year, a new generation was obtained after selection and mating of individuals belonging to the previous generation. The choice of breeders to originate the subsequent generations involved three phases: first, an individual body size selection of about 35 thousands of post-larvae (~PL21) followed by a second individual body size selection of 8-12 thousands young shrimps (~80 days old) also, free of necrosis and displaying perfect rostrum and antenna, and finally a third individual body size selection of 4000 adults (~240 days old) that are free of necrosis, displaying perfect antenna and rostrum and adequate pigmentation of gills, uropods and spermatophores. After this latter selection, 140 pairs, free of necrosis and other abnormalities, were transferred to maturation tanks and only egged females were kept at collective spawning tanks. The material sampling for genetic analysis was initiated at 1999. During the following years, samples of pleopods from shrimps belonging to generations F<sub>5</sub> ( $n=20$ ), F<sub>6</sub> ( $n=20$ ), F<sub>7</sub> ( $n=20$ ), F<sub>8</sub> ( $n=20$ ), and F<sub>9</sub> ( $n=20$ ) were collected. The tissue samples were fixed in 1 ml of 95% ethanol and kept at -20°C.

### Molecular analysis

Genomic DNA extraction followed phenol:chloroform:isoamyl protocol (Sambrook et al. 1989). Ready-To-Go™ RAPD Analysis Beads Kit (Amersham Pharmacia Biotech) was used for amplification reactions in a PTC-100™ thermal cycler (MJ Research). To each bead containing thermostable polymerases (AmpliTaq™ DNA polymerase and Stoffel fragment), 0.4 mM of each dNTP, 2.5 µg BSA, 3 mM MgCl<sub>2</sub>, 30 mM KCl, and 10 mM Tris (pH 8.3), 50 ng of template DNA, 25 pmol of a single RAPD primer and milliQ water to a 25 µl final reaction volume were added. PCR was programmed with a first cycle at 92°C for 4 min, followed by 35 cycles at 92°C for 2 min, 36°C for 1 min and 30 seconds and 72°C for 2 min with a final extension at 72°C for 3 min. The PCR products were run on agarose gel at 1.5% (for 3 h at 100 V) using 1X TBE buffer (Tris base, boric acid, EDTA) containing 0.5 µg/ml of ethidium bromide. The gels were visualized under UV light and a digital imaging system for electrophoresis gel

documentation and analysis (Kodak Digital Science™ EDAS 290) was used. From the 38 arbitrary primers tested, only those with high reproducible amplification pattern were used for statistical analyses. After the selection of a set of six primers, the amplification profiles were established and new repeatability tests were carried out for each primer. The most representative loci were identified and used in statistical analyses.

### Statistical analyses

In the RAPD analysis, dominance of alleles for the presumptive phenotypes, Mendelian segregation and populations under Hardy-Weinberg equilibrium, were assumed (D'Amato and Corach, 1996). A data matrix based on the presence (1) or absence (0) of a band for each selected locus was made involving all sampled individuals. A single- and multi-population descriptive statistics analysis for diploid data using dominant markers was performed by using the software Popgene version 1.31 (Yeh et al., 1999). Allele frequency difference among generations was calculated based on chi-square test was determined by studying multilocus genetic structure differences amongst strains. Percentage of polymorphic loci, gene diversity (Nei, 1973) and unbiased measure of genetic distance (Nei, 1972) were calculated. The Jaccard's similarity coefficient (Jaccard, 1901), which ignores the absence of bands as an indicator of similarity, was estimated by using the software NTSYS-pc version 1.8 (Rohlf, 1993). Jaccard's coefficient divergences were tested by Mann Whitney and Kruskal Wallis test using BioEstat 2.0 software program (Ayres et al., 2000).

## RESULTS

Sixty-four loci were selected for analysis and 87.5% were polymorphic. The F<sub>5</sub> and F<sub>9</sub> generations showed the largest and smallest number of polymorphic bands, respectively (Table 1). The Jaccard's similarity coefficients calculated for the sampled stocks revealed a remarkable increasing of genetic similarity levels throughout the five analyzed generations ( $P = 0.026$ ) when values from F<sub>5</sub> and F<sub>9</sub> generations were compared. Analyses of genetic diversity have also suggested a higher genetic homogeneity amongst individuals from the latter generations. The mean heterozygosity declined from 0.267 at F<sub>5</sub> generation to 0.215 at F<sub>9</sub> generation (Table 2). Analyses of gene frequencies demonstrated significant differences ( $P < 0.05$ ) at most loci, as determined by homogeneity tests (Table 3). The fragments P2-a<sub>1</sub>, P2-c<sub>3</sub>, P3-j<sub>10</sub>, P4-i<sub>9</sub>, P5-e<sub>5</sub>, P5-o<sub>15</sub>, P6-i<sub>9</sub>, and P6-a<sub>1</sub> demonstrated an increasing of their frequencies along the five generations, leading to a higher degree of homogeneity amongst individuals from latter generations. The P3-i<sub>9</sub> and P4-d<sub>4</sub> displayed progressive declined frequencies, while the fragment P4-b<sub>2</sub>, present in F<sub>5</sub>, F<sub>6</sub> and F<sub>7</sub>, was absent in F<sub>8</sub> and F<sub>9</sub> generations. Other loci also underwent to a drastic frequency reduction at F<sub>9</sub> generation, becoming present in less than 8% of individuals, for instance P4-f<sub>6</sub> and P5-b<sub>2</sub>. In contrast, two fragments, named P2-b<sub>2</sub> and P3-k<sub>11</sub>, absent from F<sub>7</sub> generation, appeared again in the F<sub>8</sub> and

**Table 1.** Nucleotide composition, number of bands and polymorphic bands per generation of each primer used in the RAPD reactions.

Oligonucleotide	Nucleotide Sequence	Number of bands	Number of polymorphic bands per generation				
			F5	F6	F7	F8	F9
Primer 1	5' GGTGCGGGAA 3'	11	06	06	06	05	05
Primer 2	5' GTTTCGCTCC 3'	07	06	05	05	05	03
Primer 3	5' GTAGACCCGT 3'	12	10	11	09	11	11
Primer 4	5' AAGAGCCCGT 3'	10	05	05	06	06	04
Primer 5	5' AACGCGCAAC 3'	15	14	13	10	10	09
Primer 6	5' CCCGTCAGCA 3'	09	08	06	07	07	03
<b>Total</b>		<b>64</b>	<b>49</b>	<b>46</b>	<b>43</b>	<b>44</b>	<b>35</b>

**Table 2.** Jaccard's genetic similarity coefficient, sample size, polymorphic loci percentage, Nei's genetic diversity (h) and the standard deviation (sd) determined for each generation (F<sub>5</sub> to F<sub>9</sub>), based on the results obtained with primers 1, 2, 3, 4, 5, and 6 in RAPD reactions.

Stock	Sample	Polymorphic loci	Jaccard's similarity	Nei genetic
F <sub>5</sub>	20	76.56%	0.621	0.267±0.187
F <sub>6</sub>	20	71.88%	0.662	0.259±0.188
F <sub>7</sub>	20	67.19%	0.701	0.242±0.198
F <sub>8</sub>	20	68.75%	0.749	0.253±0.196
F <sub>9</sub>	20	54.69%	0.795	0.216±0.211

**Table 3.** Allele frequency and homogeneity test based on the determination of chi-square ( $\chi^2$ ) and error probability ( $p$ ), where  $P < 0.05$  and  $df=4$ .

Fragments	Locus	Size (pb)	Allele Frequency (f)					Homogeneity Test	
			F <sub>5</sub>	F <sub>6</sub>	F <sub>7</sub>	F <sub>8</sub>	F <sub>9</sub>	$\chi^2$	$p$
	P1-a <sub>1</sub>	700	0.106	0.134	0.225	0.163	0.293	3.014	0.555
	P1-b <sub>2</sub>	670	0.329	0.194	0.225	0.408	0.553	7.517	0.111
	P1-c <sub>3</sub>	650	1.000	1.000	1.000	1.000	1.000	0.000	1.000
	P1-d <sub>4</sub>	560	1.000	1.000	1.000	1.000	1.000	0.000	1.000
	P1-e <sub>5</sub>	530	1.000	1.000	1.000	1.000	1.000	0.000	1.000
	P1-f <sub>6</sub>	500	0.193	0.780	0.408	1.000	0.500	41.459	0.000 <sup>S</sup>
	P1-g <sub>7</sub>	430	0.258	0.194	0.513	0.613	0.553	20.761	0.000 <sup>S</sup>
	P1-h <sub>8</sub>	410	0.225	0.194	0.780	0.163	0.776	30.366	0.000 <sup>S</sup>
	P1-i <sub>9</sub>	400	0.553	0.776	0.684	0.776	1.000	11.647	0.200 <sup>S</sup>
	P1-j <sub>10</sub>	360	1.000	1.000	1.000	1.000	1.000	0.000	1.000
	P1-k <sub>11</sub>	300	1.000	1.000	1.000	1.000	1.000	0.000	1.000
	P2-a <sub>1</sub>	710	0.777	0.776	0.776	0.776	1.000	5.446	0.244
	P2-b <sub>2</sub>	700	0.780	0.258	0.000	0.025	0.000	14.101	0.007 <sup>S</sup>
	P2-c <sub>3</sub>	660	0.776	1.000	1.000	1.000	1.000	18.726	0.001 <sup>S</sup>
	P2-d <sub>4</sub>	640	0.258	0.106	0.329	0.134	0.163	4.368	0.358
	P2-e <sub>5</sub>	550	1.000	1.000	0.776	1.000	1.000	8.726	0.001 <sup>S</sup>

Table 3. contd.


P2-f <sub>6</sub>	530	0.258	0.293	0.367	0.293	0.452	2.180	0.703 <sup>S</sup>
P2-g <sub>7</sub>	500	0.452	0.408	0.408	0.776	0.225	12.937	0.012 <sup>S</sup>
P3-a <sub>1</sub>	900	0.258	0.293	0.163	0.134	0.258	2.188	0.701 <sup>S</sup>
P3-b <sub>2</sub>	850	0.780	0.163	0.613	0.051	0.293	22.984	0.000 <sup>S</sup>
P3-c <sub>3</sub>	700	0.293	0.225	0.134	0.329	0.194	2.694	0.610
P3-d <sub>4</sub>	660	0.134	0.163	0.134	0.025	0.329	7.260	0.123 <sup>S</sup>
P3-e <sub>5</sub>	650	0.194	0.513	0.513	0.553	0.293	19.666	0.001 <sup>S</sup>
P3-f <sub>6</sub>	640	0.258	0.106	0.163	0.293	0.553	11.942	0.018 <sup>S</sup>
P3-g <sub>7</sub>	510	0.452	0.258	0.163	0.258	0.684	14.926	0.005 <sup>S</sup>
P3-h <sub>8</sub>	500	0.258	0.500	0.329	1.000	0.553	26.998	0.000 <sup>S</sup>
P3-i <sub>9</sub>	450	1.000	1.000	1.000	0.776	0.684	19.046	0.001 <sup>S</sup>
P3-j <sub>10</sub>	410	0.225	0.408	0.452	0.684	0.776	15.670	0.003 <sup>S</sup>
P3-k <sub>11</sub>	380	0.225	0.163	0.000	0.613	0.684	31.546	0.000 <sup>S</sup>
P3-l <sub>12</sub>	340	1.000	0.776	1.000	0.776	1.000	14.734	0.005 <sup>S</sup>
P4-a <sub>1</sub>	900	0.452	0.684	0.613	0.776	0.613	4.833	0.305 <sup>S</sup>
P4-b <sub>2</sub>	840	0.780	0.500	0.500	0.000	0.000	32.362	0.000 <sup>S</sup>
P4-c <sub>3</sub>	600	0.408	0.776	0.776	0.684	0.776	9.403	0.052 <sup>S</sup>
P4-d <sub>4</sub>	520	1.000	1.000	0.776	0.776	0.776	10.330	0.035 <sup>S</sup>
P4-e <sub>5</sub>	430	1.000	1.000	0.776	0.776	1.000	14.734	0.005 <sup>S</sup>
P4-f <sub>6</sub>	400	0.253	0.134	0.106	0.134	0.078	1.922	0.750 <sup>S</sup>
P4-g <sub>7</sub>	350	1.000	0.776	1.000	0.684	1.000	19.046	0.001 <sup>S</sup>
P4-h <sub>8</sub>	290	1.000	1.000	1.000	1.000	1.000	0.000	1.000 <sup>S</sup>
P4-i <sub>9</sub>	270	0.776	1.000	1.000	1.000	1.000	18.726	0.000 <sup>S</sup>
P4-j <sub>10</sub>	220	1.000	1.000	1.000	1.000	1.000	0.000	1.000 <sup>S</sup>
P5-a <sub>1</sub>	850	0.613	0.500	1.000	1.000	1.000	33.232	0.000 <sup>S</sup>
P5-b <sub>2</sub>	800	0.258	0.293	0.258	0.329	0.078	4.084	0.395 <sup>S</sup>
P5-c <sub>3</sub>	700	0.258	0.293	0.258	0.163	1.000	39.135	0.000 <sup>S</sup>
P5-d <sub>4</sub>	660	0.367	0.553	1.000	0.500	0.452	20.032	0.000 <sup>S</sup>
P5-e <sub>5</sub>	620	0.258	0.553	0.553	1.000	1.000	37.672	0.000 <sup>S</sup>
P5-f <sub>6</sub>	540	0.553	0.780	0.134	0.194	0.134	17.146	0.002 <sup>S</sup>
P5-g <sub>7</sub>	510	0.780	0.613	0.613	0.776	0.367	23.757	0.000 <sup>S</sup>
P5-h <sub>8</sub>	500	0.253	0.131	0.513	0.367	0.452	18.153	0.000 <sup>S</sup>
P5-i <sub>9</sub>	490	0.106	0.452	0.293	0.452	1.000	35.843	0.000 <sup>S</sup>
P5-j <sub>10</sub>	470	0.293	0.408	0.323	0.452	0.106	6.651	0.155 <sup>S</sup>
P5-k <sub>11</sub>	430	1.000	1.000	1.000	1.000	1.000	0.000	1.000 <sup>S</sup>
P5-l <sub>12</sub>	390	0.613	0.776	0.684	0.408	0.684	6.593	0.159 <sup>S</sup>
P5-m <sub>13</sub>	360	0.134	0.225	0.500	0.293	0.293	7.069	0.132 <sup>S</sup>
P5-n <sub>14</sub>	310	0.500	0.408	1.000	1.000	1.000	31.368	0.000 <sup>S</sup>
P5-o <sub>15</sub>	290	0.683	1.000	1.000	1.000	1.000	27.006	0.000 <sup>S</sup>
P6-a <sub>1</sub>	800	0.612	1.000	1.000	1.000	1.000	33.585	0.000 <sup>S</sup>
P6-b <sub>2</sub>	660	0.293	0.613	0.367	0.684	1.000	26.084	0.000 <sup>S</sup>
P6-c <sub>3</sub>	630	0.163	0.225	0.134	0.408	1.000	10.044	0.040 <sup>S</sup>
P6-d <sub>4</sub>	600	0.684	1.000	0.684	1.000	1.000	21.721	0.000 <sup>S</sup>
P6-e <sub>5</sub>	530	0.452	0.134	0.613	0.776	1.000	35.7032	0.000 <sup>S</sup>
P6-f <sub>6</sub>	500	0.613	0.367	0.452	0.684	1.000	20.471	0.000 <sup>S</sup>
P6-g <sub>7</sub>	470	1.000	1.000	1.000	0.776	1.000	18.726	0.001 <sup>S</sup>
P6-h <sub>8</sub>	460	0.513	0.293	0.106	0.258	0.258	5.931	0.204 <sup>S</sup>
P6-i <sub>9</sub>	420	0.106	0.258	0.293	0.329	0.684	16.360	0.003 <sup>S</sup>

S: significantly difference, P<0.05.

**Table 4.** Genetic distance based on the Nei's unbiased measure.

Stock	Nei's Unbiased Measure of Genetic Distance				
	F <sub>5</sub>	F <sub>6</sub>	F <sub>7</sub>	F <sub>8</sub>	F <sub>9</sub>
F <sub>5</sub>					
F <sub>6</sub>	0.044	*****			
F <sub>7</sub>	0.060	0.038	*****		
F <sub>8</sub>	0.112	0.084	0.081	*****	
F <sub>9</sub>	0.139	0.116	0.122	0.077	*****

F<sub>9</sub> generations, what probably can reflect a sampling artifact. The results obtained from genetic diversity analysis demonstrated differentiated levels of genetic distance among the stocks (Table 4).



**Figure 1.** Jaccard's similarity coefficient against number of generations in the broodstock line studied. Correlation ( $R^2$ ) = 0.9845.

## DISCUSSION

Allele frequency distribution along the five surveyed generations showed significant differences at most of the analyzed loci likely related to genetic drift and inbreeding. A putative random founder effect during the selection process of broodstock can lead to an effective loss of alleles in the following generations, just as verified for the fragment P4-b<sub>2</sub> absent in the generations F<sub>8</sub> and F<sub>9</sub>. In contrast, several other alleles were only detected in the latter generations and could account for inbreeding effects, increasing their frequencies along the generations, although small sampling size effects could not be discarded.

Inbreeding effects should also lead to the observed decreased genetic diversity along the generations, verified by a strong positive correlation between genetic

similarity and generation time ( $R^2 = 0.9845$ ) and the genetic similarity could reach 100% near to 18-19 generation in this commercial line studied (Figure 1).

Similar findings have already been reported in other aquaculture organisms. Allozyme allele loss and heterozygosity reduction were reported when captive stocks of *L. vannamei* were compared with wild populations (Sunden and Davis, 1991). Allele losses were also revealed in reared lines of *Litopenaeus stylirostris*, and divergences observed between two studied lines indicated a probable founder effect since both stocks shared a common origin from a single wild population (Ramos-Paredes and Grijalva-Chon, 2003).

Founder effect and inbreeding have also been identified in *M. japonicus* populations, revealing a continuous reduction of allozyme polymorphism level throughout six captive generations, from 0.102 at F<sub>1</sub> to 0.039 at F<sub>6</sub> generation. An initial bottleneck which occurred on the first generation might have favored to a remarkable reduction of genetic variation levels in the following ones. Although the number of shrimp pairs used to produce the next generation ranged from 50 to 300, this study revealed that the effective number of parents contributing to each broodstock was as low as four (Sbordoni et al., 1986).

The commercial line here analyzed was formerly founded with animals originated from different countries (Panama, Ecuador, Venezuela, Costa Rica, and Mexico) and they have undergone a domestication process, currently reaching up to F<sub>9</sub> generation. Despite of this, diverse origin has contributed to a higher genetic diversification at the founder population since new gene pools were never introduced since then. The maintenance of this closed line throughout generations has favored the increasing of inbreeding and, consequently, to a higher consanguinity level. Although, thus far, there were no evidence of performance losses regarding to fecundity and growth rates in the shrimp lineage studied here which could be accounted to inbreeding depression.

The increase of the effective number of mating pairs could be a useful strategy to diminishing the reduction rate of genetic diversity within this line, and cross-breeding using shrimp of different origins could recover

part of genetic variability lost in this commercial closed line.

## ACKNOWLEDGEMENTS

This research was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Associação Brasileira de Criadores de Camarão (ABCC). The authors thank Maricultura shrimp hatchery and farm for the shrimp samples and information about the stocks studied.

## REFERENCES

- Ayres M, Ayres Jr M, Ayres DL, Santos AS (2000). Biostat 2.0: Aplicações Estatísticas nas Áreas de Ciências Biológicas e Médicas. CNPq, Brasília.
- Batalha MO, Silva AL, Souza-Filho HM, Galetti PMJ, Pigatto G, Freitas PD, Lucchese T (2002). A Maricultura no Estado de São Paulo. Sebrae, São Paulo, 297 pp.
- Bradford GE, Chapman AB, Grummer RH (1958). Effects of inbreeding selection linecrossing and topcrossing in swine. 1. Inbreeding and selection. *Animal Science* 17: 426-440.
- D'Amato ME, Corach D (1996). Genetic diversity of populations of the freshwater shrimp *Macrobrachium borelli* (Caridae: Palaemonidae). *J Crust Biol.* 16(4): 650-655.
- Galetti Jr PM, Freitas PD (1999). A genética de camarão no Brasil. *Rev da ABCC* 1(1): 30-32.
- Gilpin ME, Soulé ME (1986). Minimum viable populations: processes of species extinction. In: Soulé ME (ed.) *Conservation Biology: The science of scarcity and diversity*. Sinauer Associates, Sunderland, MA. pp. 19-34.
- Jaccard P (1901). Étude comparative de la distribution florale dans une portion des Alpes et des Jura. *Bull. Soc. Vaudoise Sci. Nat.* 37: 547-579.
- Nei M (1972). Genetic distance between population. *Am. Nat.* 106: 283-292.
- Nei M (1973). The theory and estimation of genetic distance. In: Morton NE (ed.) *Genetic structure of populations*. University of Hawaii, Honolulu, pp. 45-54.
- Ramos-Paredes J, Grijalva-Chon JM (2003). Allozyme genetic analysis in hatchery strains and wild blue shrimp, *Penaeus (Litopenaeus) stylirostris* (Stimpson), from the Gulf of California. *Aquaculture Research* 34: 221-234.
- Rocha IP, Rodrigues J, Amorim L (2004). A carcinicultura brasileira em 2003. *Rev da ABCC* 6 (1): 30-36.
- Rohlf FJ (1993). Numerical taxonomy and multivariate analysis system, NTSYS-pc, v 1.80 Applied Biostatistics Inc., New York.
- Sambrook J, Fritsch EF, Maniatis T (1989). *Molecular cloning: A Laboratory Manual* 2nd ed. Cold Spring Harbor, New York.
- Sbordoni V, De Matthaeis E, Cobolli -Sbordoni M, La Rosa G, Mattoccia M (1986). Bottleneck effects and depression of genetic variability in hatchery stocks of *Penaeus japonicus* (Crustacea, Decapoda). *Aquaculture* 57: 239-251.
- Sunden SLF, Davis SK (1991). Evaluation of genetic variation in a domestic population of *Penaeus vannamei* (Boone): A+ comparison with three natural populations. *Aquaculture* 97:131-142.
- Tave D (1993). *Genetics for fish hatchery managers*, 2nd ed. Van Nostrand-Reinhold, New York. pp. 415.
- Vrijenhoek RC (1994). Genetic diversity and fitness in small population. In: Loeschcke V, Tomiuk J, Jain SK (eds.) *Conservation Genetics*. Birkhäuser Verlag Basel, Switzerland. pp. 38-53.
- Yeh F, Yang R, Boyle T (1999). Microsoft windows-based freeware for population genetics analysis. Popgene version 1.31. <http://www.ualberta.ca/~fyeh/download.htm>.