

Full Length Research Paper

Antibacterial and antifungal screening of concentrates from six Medicinal plants gathered in Kinshasa-Democratic Republic of Congo against clinical detach pathogens

*Antonio Kabila, Augustine Tshombe and Kaumba E. B

Department of Pharmacognosy, Faculty of Pharmacy, University of Goma, Democratic Republic of the Congo.

Accepted 20 February, 20119

Results from the *in vitro* evaluation of the antibacterial and antifungal activities of six plant extracts indicated that the aqueous and the methanol extracts of *Acalypha wilkesiana* leaves and *Ageratum conyzoides* leaves exhibited good and efficient antifungal activity against *Candida albicans* with minimum inhibitory concentration (MIC) of 31.25 µg/ml and minimum fungicidal concentration (MFC) of 62.5 µg/ml while the methanol extract from *Pentaclethra macrophylla* inhibited the yeast growth with minimum inhibitory concentration (MIC) and minimum fungicidal concentration (MFC) values of 62.5 and 125 µg/ml respectively. The aqueous extracts from *Acalypha wilkesiana*, *Ageratum conyzoides*, *Bucchozia tholoniana* seeds, *Gulboutia demeusei* root and the aqueous and methanol extracts from *Pentaclethra macrophylla* stem bark showed good antibacterial activity against *Escherichia coli*, *Klebsiella oxytoca*, *Proteus mirabilis*, *Salmonella thyphimurium*, *Stahylococcus aureus* and *Shigella flexneri* with minimum inhibitory concentration (MIC) of 62.5 µg/ml and minimum bactericidal concentration (MBC) of 125 µg/ml according to the case. These results partly support and justify the traditional use of these plant extracts for treating infections in traditional medicine.

Key words: Plant extracts, antibacterial, antifungal, infections.

INTRODUCTION

Bacteria's are responsible for serious and various human and animal infections. For example, *Pseudomonas*

aeruginosa is the most common bacteria in clinics with significant percentage of acquired infections (Abu-

*Corresponding author. E-mail: paper_submit@gmail.com

Shanab et al., 2004). *Sataphylococcus aureus* causes skin lesions which are often superficial, localized abscesses and food contamination or poisoning (Lotifpour et al., 2008). Intestinal disorders are caused by bacteria such as *Vibrio cholerae*, *Escherichia dysenteriae*, *Escherichia coli* and *Salmonella* species especially diarrhoea as a major cause of mortality and morbidity in developing countries (Chitemerere and Makangangama, 2011). Long before mankind the existence of *microbes* was discovered, the idea that some medicinal plants had antibacterial and antifungal potential since they contain active constituents belonging to different phytochemical groups, was well accepted (Rios; Recio, 2005). Thus, the therapeutic properties of various medicinal plants have been known to treat particularly human and animal infectious diseases as it is estimated between 60 to 90% of the population in developing countries turn to traditional medicine and consider it to be a normal part of its primary healthcare (WHO, 2002).

The rapid spread of bacteria and fungi expressing multidrug resistance (MDR) has encouraged the discovery of new antibacterial and antifungal agents. The trend use in alternative and complementary healthcare has prompted scientists to investigate various biological activities of medicinal plants mainly according to their uses in traditional medicine to prove their effectiveness and safety to support and justify their traditional uses. Crude extracts prepared from medicinal plants as infusion, decoction, macerate, tincture or powder are traditionally used by the population to treat various diseases particularly infections from different origins. Although their efficacy and mechanism of action have not been scientifically elucidated in most cases, these different medicinal preparations often give beneficial response to human and animal due to the presence of active compounds (Barnes et al., 2007).

In this present study, six medicinal plants belonging to different botanical families were selected and evaluated *in vitro* for their potential antibacterial and antifungal properties. All selected plant parts are used in traditional medicine in Kinshasa-Democratic Republic of Congo (DR Congo) to treat various ailments such as diarrhoea, wounds, burns and skin infections (Kerharo and Adam, 1974), (Oliver Bever, 1986), (Kambu, 1990), (Neuwinger, 2000). To treat wounds and burns, powder or decoction from the plant part is applied and this process gives good results as indicated by the acceleration of the rapid healing of affected part of the body and the decoction is employed to treat bacterial diarrhoea. For this study, the aqueous and methanol extracts from the selected plant parts were tested against 9 clinical isolate pathogens and 1 yeast *in vitro*. Their minimum inhibitory concentrations were determined by dilution method when compared to the reference antibacterial and antifungal products.

MATERIALS AND METHODS

Selected Plants

They include leaves of *Acalypha wilkensis* Mull. Arg. (Euphorbiaceae), leaves of *Ageratum conyzoides* L. (Asteraceae), seeds of *Buchholzia tholoniana* Hua (Capparidaceae), root of *Gulbournia demeusei* (Harms) J. Leonard (Fabaceae), stem bark of *Pentaclethra macrophylla* Benth (Mimosaceae) and leaves of *Syzygium guineense* (Wild.) DC. var. *guineense* (Myrtaceae) collected in Kinshasa-Democratic Republic of Congo (DR Congo) in April 2011. All plants were identified by Mr Nlandu Lukebiako, B. of the Institute National d'Etudes et de Recherches en Agronomie, Department of Biology, Faculty of Sciences, University of Kinshasa. A voucher specimen of each plant NL01042011AW, NL02042011AC, NL03042011BT, NL04042011GD, NL05042011PM and NL06042011SG for *A. wilkensis* leaves, *A. conyzoides* leaves, *B. tholoniana* seeds, *G. demeusei* root, *P. macrophylla* stem bark and *Syzygium guineense* leaves respectively has been deposited in the herbarium of this institute. All plant parts were dried at room temperature and reduced to powder.

Preparation of extracts

45 g of each dried plant material were macerated with distilled water or methanol (200 ml each) during 24 h. After, each mixture was filtered and each filtrate evaporated in vacuum yielding corresponding dried extracts (Table 1).

Phytochemical screening

This study was performed by thin-layer chromatography (TLC) on precoated silica gel plates F₂₅₄ (thickness later 0.25, mm, Merck, Germany) using different reagents and mobile phases described in the literature for the identification of major chemical groups such as alkaloids, anthraquinones, coumarins, flavonoids, terpenes and steroids. Hydrochloride acid 2M/*n*-butanol (heating with HCl 2 M for 40 min at 100°C and extracting the red color with iso-amylic alcohol, froth test and Stiasny's reagent (formol + HCl conc.) were used to detect anthocyanins, saponins and tannins respectively (Harborne, 1998).

Antibacterial and antifungal testing

Selected test microorganisms

The selected microorganisms included the yeast *C. albicans*, and the bacteria *E. coli*, *Enterobacter cloacae*, *Klebsiella pneumoniae*, *Klebsiella oxytoca*, *Proteus mirabilis*, *Shigella flexneri*, *S. thyphimurium* and *S. aureus*. The yeast and all bacteria were clinic isolated from patients diagnosed with infections in the Cliniques Universitaires du Mont-Amba/Kinshasa-DR Congo.

Culture medium

Bacteria and fungi were maintained on trypticase soy broth (TSB) plates and Sabouraud dextrose agar plates at 4°C respectively. Yeast were cultured in liquid Sabouraud dextrose medium for 48 h at 24 °C. Test inocula of fungi were prepared by harvesting matured sporulating cultures in Sabouraud agar broth. For inoculation with the yeast, homogenized mycelia cultures of two weeks old were

Table 1. Uses in traditional medicine of ethnopharmacologically selected medicinal plants

Plant	Uses
<i>Acalypha wilkesiana</i> Mull. Arg. (Euphorbiaceae)	The leaves are used to treat fungal skin diseases and function infections, pityriasis vesicolor, superficial mycoses. Associated to <i>Ocimum gratissimum</i> against methicillin resistant bacteria, to combat constipation, naso-pharyngeal infections, pain-killers, applied externally as an antiseptic to ulcers and wounds, antidiarrhoeal agent, used as antibiotic, bacteriostatic and fungistatic. The plant possess anticancer, antibacterial, antifungal, antidiabetic, analgesic, antiparasitic and antihypertensive properties and so on (Neuwinger,, 2000; Lim et al., 2013: Royal Botanic Gardens Kew, 2014).
<i>Ageratum conyzoides</i> L. (Asteraceae)	The grain cannot be used if it is contaminated due to its toxicity. Leaves and entire plant are used to treat colic, colds and fever, diarrhea, rheumatism, spasms, headache, pneumonia, wounds, burns, and as a tonic. They are also recommended to treat burns, wounds infectious conditions and bacterial infections. The leaf sap is used to treat conjunctivitis, as galactagogue, liver and stomach problems, otitis, and so on (Anonymous, 2014).
<i>Bucchozia tholoniana</i> Hua (Capparaceae)	The seed is used to treat inflammation, gynecological infections, migraine, cough and stomach pains (Oliver Bever, 1986).
<i>Gulbourtia demeusei</i> (Harms) J. Léonard (Fabaceae)	The stem bark is employed to treat otitis, wound, skin infections and diarrhea (Oliver Bever, 1986)
<i>Pentacletra macrophylla</i> Benth. (Euphorbiaceae)	The bark is employed as a decoction to treat abdominal pains, dysmenorrhoeal, urogenital infections, gonorrhoea, stomach pains, dysentery, hernia, as anthelmintic, The ripe seeds and stem bark powder are applied externally to treat wounds, and burns, Extracts of the leaf, stem bark, seeds and seed pulp have antiinflammatory and antihelmintic properties. They are used to treat gonorrhoea and convulsions, and as analgesic. The root and stem barks are used as a laxative, enema against dysentery and liniment against itch. It is also used to treat stomach pains, wounds, burns asthma, abscess, heart pains, diarrhoea and ulcerations, and so on (Kerharo and Adam, 1974: Kambu, 1990; Neuwinger, 200).
<i>Syzygium guineense</i> (Wild.) DC. (Myrtaceae)	The stem bark or leaf have febrifuge, anthelmintic and purgative properties They also are used to treat menstrual cycle, stomach pains, colics, dysentery, rheumatism pains, malnutrition, delirium, naso-pharyngeal affections, pain killers, pulmonary troubles and diarrhoea. The leaf powder is a healing agent. An infusion of bark and roots soaked in hot water is used as a purgative, the bark can be toxic and deaths from its use have been recorded. The root is soaked in water for drinking and bathing to treat epilepsy, Root extracts are used as taeniocide. Bark decoction is employed against stomach-ache, diarrhoea and malaria, as a mildly laxative and applied in draught or in baths as a tonic. Its infusion is taken against coughs, asthma, throat problems and intercostals pains. The bark is used for the treatment of snakebites and so on (Kerharo and Adam, 1974; Maroyi, 2008; Kupumba, 2014).

Kerharo and Adam, 1974, Kambu, 1990, Neuwinger, 2000, Anonymous,2014; Maroyi, 2008. Lim et al., 2013; Kupumba, 2014, Royal Botanic Gardens Kew, 2014.

used (Vanden Berge and Vlietinck, 1991; Cimanga et al., 1998).

Antibacterial and antifungal testing

The antimicrobial and antifungal tests extracts used against bacteria and yeast respectively were performed by the microtiter plate dilution method. For this technique, 2 mg of each extracts was dissolved in dimethylsulfoxide (DMSO), water and trypticase soy broth (TSB), final volume 2 ml (concentration of DMSO < 1%) to have a final concentration of 1 mg/ml. These stock solutions were repeatedly diluted in two fold with TSB or Sabouraud medium to obtain a series of test concentrations samples from 500 to 10 µg/ml.

An inoculum consisting of about 10^5 microorganisms/ml TSB or 10^6 yeast /ml Sabouraud medium was incubated overnight at 37°C for 24 h. A 1/1000 dilution of each suspension was prepared with the corresponding medium. Thereafter, 1 ml of the test sample was added. Each vertical column contained 100 µl TSB and bacteria or 100 µl Sabouraud medium and yeast without test sample was in control for the normal growth of bacteria or yeast. The plates were incubated at 37°C in humidified atmosphere for 24 h. The inhibition of bacterial and yeast growth was evaluated by comparing it with normal bacterial or yeast growth in control holes prepared without test samples. The MIC was determined as the lowest concentration of the sample that completely inhibited macroscopic growth of bacteria or yeast. To determine the minimum bactericidal or

Table 2. Amount plant extracts (g) from 45 g of plant material

Plant species	Part used	Aqueous extract (%)	Methanol extract (%)
<i>Acalypha wilkesiana</i> Mull. Arg	L	1.94 (4.31)	3.94 (8.76)
<i>Ageratum conyzoides</i> L.	L	4.5 (10)	4.4(9.98)
<i>Buchholzia tholoniana</i> Hua	F	1.37 (3.04)	2.70 (6.00)
<i>Guilourtia demessii</i> (Hams) J. Leonard	R	3.07 (6.82)	7.27(16.16)
<i>Pentaclethra macrophylla</i> Benth.	Sb	0.8 (1.78)	6.25(13.89)
<i>Syzygium guineense</i> (Wild) D.C.	L	1.44 (3.20)	7.20(16.61)

L: leaf, R: root, Sb: stem bark

Table 3. Phytochemical screening of the aqueous extracts of selected plant parts

Chemical groups/Plant extract	AC	AW	BT	GD	PM	SG
Alkaloids	++	++	-	++	++	-
Anthraquinones	-	-	+	-	+	-
Anthocyanins	-	-	-	-	-	-
Coumarins	++	-	+	-	-	-
Flavonoids	+	++	+	+	+	+
Tannins	+	+	-	-	++	++
Terpenes/steroids	++	++	++	++	++	++
Saponins	-	-	-	+	+	+
Sugars	-	-	+	+	+	+

These chemical groups were also detected in each corresponding methanol extract. AC: *Ageratum conyzoides*, AW: *Acalypha wilkesiana*, BT: *Buchholzia tholoniana*, GB: *Guibourtia demeusei*, PM: *Pentaclethra macrophylla*, SG: *Syzygium guineense*.

fungicidal concentration (MBC or MFC), the two lowest concentrations which inhibited bacterial or yeast growth were plated out on a nutrient agar and incubated at 37°C for 24 h. Results were evaluated by comparing them with the control holes containing bacteria or yeast without test sample (Vanden Berghe and Vlietinck, 1991; Cimanga et al., 1998).

RESULTS AND DISCUSSION

The type and level of different biological activities exhibited by any plant extract is dependent on many factors such as geographical area, time of collection, soil conditions, harvesting time, moisture content, drying method, storage conditions and post-harvest process (Wendakoon; Calderon, 2012). Also, the relatively high temperature and humidity can be generated during tissue grinding and can denature chemical constituents, the extractive solvent can also affect the level and composition of secondary active metabolites extracted. It is well known that the bacteriostatic effect of an antibiotic is sufficient to treat infections in correlation with the host's defense such as cellular and humoral systems while severe infections necessitate a rigorously bactericidal or

fungicidal agent administered together which may produce a synergistic effect (Hardman and Limbird, 2001; Wheels, 2008). In the present study, Table 1 reports some traditional uses of the six ethnopharmacologically selected medicinal plants while Table 2 shows the amounts of the aqueous and methanol extracts of each treated plant part. The six medicinal plants chosen for this study are commonly used for the treating diverse infections in traditional medicine in Kinshasa according to the daily practices of interviewed traditional healers and the literature data. Some of them are known to produce bioactive products (Djoukeng et al., 2005). Table 3 reports results from the phytochemical screening of the aqueous extract of all selected plant parts. The chemical composition of each methanol extract was similar to that of the corresponding aqueous extract. These results revealed the presence of alkaloids in *Guibourtia demeusei*, *Pentaclethra macrophylla* and *Syzygium guineense* extracts, flavonoids, steroids and terpenes in all plant extracts, tannins in *Ageratum conyzoides*, *Acalypha wilkesiana*, *Pentaclethra macrophylla* and *Syzygium guineense*, saponins in *Guibourtia demeusei*, *Pentaclethra macrophylla* and *Syzygium guineense*

extracts and sugars in *Buccholzia tholoniana*, *Guibourtia demeusei* and *Pentaclethra macrophylla*. Coumarins were only detected in *Buccholzia tholoniana* while anthocyanins were not found to be present in all extracts in our experimental conditions.

The dilution method used in this study has been currently employed in other previous investigations (Kambu et al., 1990; Vlietinck et al., 1995; Fabry et al., 1998; Cimnaga et al., 1998; David et al., 2009; Singla et al., 2011; Djeussi et al., 2013) and is recommended as a good method for determining the relative potency of crude extracts and establishing their antibacterial and antifungal spectrum as it facilitates the use of different strains and to determine their MIC and MBC or MFC. It concerns polar, apolar and complex extracts in testing to establish their real potency by determining their MIC and MBC or MFC (Rios; Recio, 2005). Beside the dilution procedure, the disc diffusion method is also frequently used with high test concentrations from 1 to 100 mg to evaluate antibacterial and antifungal activity of plant extracts, it is only a qualitative technique since the MIC and MBC are not determined (Kelmanson et al., 2000; Samy and Ignacimuthu, 2000; Islam et al., 2008; David et al., 2009; Jamuna Bai et al., 2011; Chitemerere et al., 2011; Muhamed Muback et al., 2011; Neeraj and Madivi, 2011; Debajit et al., 2012; Israr et al., 2012; Sakunpak and Panichayupakaranant, 2012; Sharmeen et al., 2012; Tabasum et al., 2013; Gupta et al., 2013). To avoid this inconvenience, both techniques are used at the same time (Jagessar et al., 2008; Oskay et al., 2009; Chitemerere and Mukanganyama, 2011; Kondo et al., 2010; Shinkafi and Dauda, 2013; Raghavendra and Mahadevan, 2011; Dugler and Dugler, 2012; Sadeghi-Nejad and Azish, 2013) Antibacterial and antifungal activities of plant extracts or other compounds is largely influenced by the susceptibility testing method used (Vanden Berghe and Vlietinck, 1991).

For the present study, the following criteria were adopted to appreciate the level of evaluated activities: MIC, MBC or MFC 10 µg/ml: pronounced activity, 10 MIC, MBC or MFC < 100 µg/ml: good activity, 100 MIC, MBC or MFC 125 µg/ml: moderate activity, 125 < MIC, MBC or MFC 250 µg/ml: low activity, 250 < MIC, MBC or MFC 500 µg/ml: very low activity, MIC, MBC or MFC > 500 µg/ml: inactive. Results presented in Table 4 shows the MICs of tested plant extracts. They show that the aqueous and the methanol extracts from all selected medicinal plants exhibited antibacterial (bacteriostatic) and antifungal (fungistatic) activity at different extents.

When tested against the yeast *C. albicans*, the most active extracts with good activity were the aqueous extracts from *A. wilkesiana* leaves and *A. conyzoides* leaves inhibiting the yeast growth activity with MIC value of 31.25 µg/ml. The methanol extracts from *A. wilkesiana*,

A. conyzoides and *Pentaclethra macrophylla* stem bark also presented good antifungal activity with MIC value of 62.5 µg/ml. The aqueous extracts of *Buccholzia tholoniana* seeds and *P. macrophylla*, and the aqueous and methanol extract of *Syzygium guineense* leaves showed moderate activity with MIC value of 125 µg/ml. The remaining samples displayed low, very low antifungal activity or were inactive against the selected yeast (Table 4). The observed effect of these plant extracts against *C. albicans* was fungistatic at different degrees. Results in Table 4 also show the effect of these extracts against 9 clinical isolate pathogens. They indicated that the aqueous from *A. conyzoides* exhibited good antibacterial activity (bacteriostatic effect) against *E. coli* with MIC value of 62.5 µg/ml. The same minimum inhibitory concentration was also presented by the methanol extracts from *P. macrophylla* and *S. guineense* against

Enterobacter species, the aqueous extracts from *A. wilkesiana* and *P. macrophylla*, and the methanol extract from *P. macrophylla* against *Klebsiella oxytoca*, the aqueous extracts from *A. wilkesiana*, *G. demeusei*, *P. macrophylla* and the methanol extract of this last plant against *Proteus mirabilis*, the aqueous extracts from *A. conyzoides* and *P. macrophylla* against *Salmonella thyphimurium*, the aqueous extract of *B. tholoniana* against *S. aureus*. The remaining samples were found to display moderate activity (MIC = 125 µg/ml), low (125 < MIC 250 µg/ml), very low (250 < MIC 500 µg/ml) or were inactive (MIC > 500 µg/ml) against the selected clinical isolate bacteria (Table 4).

Table 5 shows the minimum bactericidal (MBCs) and fungicidal (MFCs) concentrations of the selected plants extracts. Results indicated that the aqueous extracts from *A. wilkesiana* and *A. conyzoides*, and the methanol extract from *P. macrophylla* presented good fungicidal activity with MBC value of 62.5 µg/ml against *C. albicans*. The aqueous and methanol extract from *P. macrophylla* exhibited good activity against *K. oxytoca* (MBC = 62.5 µg/ml). The same MBC was also shown by the aqueous extracts of *A. wilkesiana*, *G. tholoniana*, the aqueous and methanol extract of *P. macrophylla* against *P. mirabilis*, the aqueous extract of *A. conyzoides* and *P. macrophylla* against *S. thyphimurium*. The remaining plant extracts showed moderate, low and very low bactericidal and fungicidal activities or were devoid with these effects according to the case (Table 5). The bacteriostatic and bactericidal effect of the aqueous extracts of *A. conyzoides*, *A. wilkesiana* and *G. demeusei*, and the methanol extract of *P. macrophylla* was higher compared to that of norfloxacin, used an antibiotic reference product against some bacteria such as *E. coli*, *E. cloacae*, *K. oxytoca*, *P. mirabilis*, *S. thyphimurium*, *S. aureus* and *S. flexneri* according to the tested bacteria. The activity of the remaining extracts was sometime similar or lower

Table 4. Antibacterial and antifungal activities of plant extracts (MIC, µg/ml)

Extract	Microorganism								
	<i>C. a.</i>	<i>Es. c.</i>	<i>En. c.</i>	<i>K. p.</i>	<i>K. o.</i>	<i>P. m.</i>	<i>S. f.</i>	<i>S. a.</i>	<i>S. t.</i>
<i>A. wilkesiana</i>									
Aqueous	31.1	125	125	250	62.5	62.5	125	250	125
Methanol	62.5	250	250	250	125	125	500	125	500
<i>A. conyzoides</i>									
Aqueous	31.5	62.5	125	250	250	125	62.5	250	125
Methanol	62.5	125	250	125	250	250	125	250	250
<i>B. thioniana</i>									
Aqueous	125	1000	500	125	250	500	250	62.5	1000
Methanol	1000	1000	500	250	500	500	250	125	250
<i>G. demeussi</i>									
Aqueous	500	125	500	250	125	62.5	125	125	62.5
Methanol	500	250	250	250	250	125	250	250	125
<i>P. macrophylla</i>									
Aqueous	125	125	125	125	62.5	62.5	62.5	125	125
Methanol	62.5	125	62.5	250	62.5	62.5	125	125	250
<i>S. guineense</i>									
Aqueous	250	500	125	500	125	125	500	500	250
Methanol	125	500	62.5	500	250	250	500	500	250
Norfloxacin	-	125	62.5	250	500	125	250	125	125
Tetracycline HCl	-	7.81	7.81	3.90	3.90	1.95	0.97	0.98	0.97
Nystatin	0.25	-	-	-	-	-	-	-	-

Legend: *C.a.*: *Candida albicans*, *Es.c.*: *Escherichia colis*, *En.c.*: *Enterobacter cloacae*, *K.p.*: *Klebsiella pneumonia*, *K.o.*: *Klebsiella oxytoca*, *P.m.*: *Proteus mirabilis*, *S.f.*: *Schigella flexneri*, *S.a.*: *Staphylococcus aureus*, *S.t.*: *Staphylococcus thyhimurium* -: no tested.

and 5). According to Bauernfeind and Petermüller (1983) and Boyd et al. (2009), MICs of norfloxacin for isolate pathogens from hospital ranged from 0.03 to 0.15 µg/ml. Results from our study showed that this antibiotic had high MICs against all clinical isolated bacteria tested in the present study. This finding suggested that these bacteria are resistant to norfloxacin. Tetracycline used also as an antibiotic reference product presented high antibacterial activity than norfloxacin and all tested extracts. However, the antifungal and fungicidal activity of all tested extracts was lower than that of nistatin used as an antifungal reference product (Tables 4 and 5).

Our results are only qualitatively in good agreement with those previously reported by Kambu et al. (1990) and Tona et al. (1999) concerning the antibacterial activity of *P. macrophylla* stem bark, the antibacterial and antifungal

activity of *A. conyzoides* leaves described by Ahmagboul et al.,(1985), Yamamoto et al., (1991), Gamesan and Krishnaraju (1995), Dayie et al., (2008) and Okwori et al. (2013) and the antibacterial and antifungal activity of *A. wilkesiana* leaves reported by Alade and Irobi (1993); Adesina et al., (2000); Gotep et al., (2010) and Haruna et al. (2013). Although these activities were carried out by the diffusion method. Gallic acid, corilagin and geraniin isolated from the leaves were reported to be responsible for the observed antibacterial activity (Adesina et al., 2000). In clinical trial on dermatitis, the water extract was found to be reasonably useful in the treatment of *eczema* and no allergy or irritation was observed in treated patients (Jekayinfa et al., 1997).

A previous investigation of the antibacterial activity of the methanol *A. conyzoides* whole plant extract was

Table 5. Antibacterial and antifungal activities of plant extracts (MBC and MFC, µg/ml).

Extract	Microorganism								
	<i>C. a.</i>	<i>Es. c.</i>	<i>En. p.</i>	<i>K. p.</i>	<i>K. o.</i>	<i>P. m.</i>	<i>S. f.</i>	<i>S. a.</i>	<i>S. t.</i>
<i>A. wilkesiana</i>									
Aqueous	62.5	250	250	500	125	62.5	125	500	250
Methanol	125	500	500	250	250	125	500	250	500
<i>A. conyzoides</i>									
Aqueous	62.5	125	250	500	250	125	62.5	250	125
Methanol	125	250	250	125	250	250	125	250	250
<i>B. tholoniana</i>									
Aqueous	250	1000	500	250	500	500	500	125	1000
Methanol	1000	1000	500	500	500	500	250	125	250
<i>G. demeussi</i>									
Aqueous	1000	250	1000	500	250	62.5	250	250	125
<i>P. macrophylla</i>									
Aqueous	125	125	125	125	62.5	62.5	62.5	125	125
Methanol	62.5	125	62.5	250	62.5	62.5	125	125	250
<i>S. guineense</i>									
Aqueous	500	1000	250	1000	125	250	500	500	500
Methanol	250	1000	125	500	250	500	500	500	500
Norfloxacin	-	125	62.5	250	500	125	250	125	125
Tetracycline	-	15.62	15.61	7.80	7.80	3.90	1.95	1.95	0.97
Nystatin	0.25	-	-	-	-	-	-	-	-

compared to that of the leaves and flowers, and the results had indicated that the two last extracts were more effective than the first one against the tested bacteria such as *S. aureus*, *E. coli* and *P. aeruginosa* (Dayie et al., 2008). The essential oil of *A. conyzoides* leaves was also reported to have antibacterial activity against a large microorganisms and yeasts by diffusion method (Osho and Adetunji, 2011). In addition, it had been reported that wounds treated with the aqueous leaves extract in combination with honey and with solcosery ointment significantly accelerate wound healing process and the rates of wounds sterility compared to wound treated with honey alone (Mustafa et al., 2005). Another study conducted by Dash and Murthy (2011) demonstrated that the methanol and aqueous extract from *A. conyzoides* leaves showed father rate of wound healing compared to

the effect of petroleum and chloroform extract from the same plant part. Among these later extracts, the chloroform extract was found to produce promising effect, but lower compared to that of the methanol and aqueous extracts, while the petroleum ether extract did not produce significant effect. In a previous study, a series of triterpene compounds were isolated from the leaves of *S. guineense* among which arjumolic acid and asiatic acid showed the most significant antibacterial activity against *E. coli*, *B. subtilis* and *S. sonnei* (Djoukeng et al., 2005).

Our results also indicated that the antibacterial and antifungal activities vary with the plant species, the plant part used and the nature of the extractive solvent (Malesh and Satish, 2008; Rios and Recio, 2005). Moreover, these reported results in sometime compared to those from previous screenings of other medicinal plants for the

same biological activities, showed good effect against Gram-positive strains while Gram-negative remained resistant (Vlietinck et al., 1995), (Rabe and van Staden, 1997). The observed biological activities in all selected plant extracts are related to the presence of some secondary metabolites such as terpenes, steroids, alkaloids, flavonoids, saponins and tannins found to be present in these crude extracts for which they are well known to possess these activities at different extents (Helberg et al., 1983; Ebi et al., 1997, Cowan, 1999; Mendonça Filho, 2006, Das, 2010).

Conclusion

These selected plant extracts which are used for the treatment of several infections in traditional medicine demonstrate an appreciable *in vitro* antibacterial and antifungal, and at some extents bactericidal and fungicidal activities. Thus there is now a scientific validation for the use of the selected plant part extracts for the treating of infectious diseases in traditional medicine.

Conflict of Interests

The author(s) have not declared any conflict of interests.

REFERENCES

- Abu-Shanab B, Adwan G, Abu-Saliya D, Jarra N, Adwan K (2004). Antibacterial activities of some plant extracts utilized in popular medicine in Palestine. *Turk. J. Biol.* 28:99-102.
- Adesina SK, Idowu O, Ogundaini AO, Oladimeji H, Olugbade TA, Onawunmi GO, Pais M (2000). Antimicrobial constituents of the leaves of *Acalypha wilkesiana* and *Acalypha hispida*. *Phytother. Res.* 14(5):371-374. doi:10.1002/1099_1573(200008)14:5<371::AID-PTR625>3.0.CO;2-F
- Ahmagboul AZ, Farroq AA, Tyagi BR (1985). Antimicrobial activity of certain Sudanese plants used in folkloric medicine: Screening for antibacterial activity, part II. *Fitoterapia.* 56:103-109.
- Alade PI, Irobi ON (1993). Antimicrobial activities of leaf extracts of *Acalypha wilkesiana*. *J. Ethnopharmacol.* 39:171-174.
- Anonymous (2014). Tropical Plant database, *Ageratum (Ageratum conyzoides)*. Raintree. www.rain-tree.com/ageratum.htm
- Barnes JI, Anderson A, Phillipson JD (2007). *Herbal Medicine*, 3rd Ed.
- Bauernfeind A, Pettermüller C (1983). In vitro activity of ciprofloxacin, norfloxacin and nalidixic acid. *Eur. J. Clin. Microbiol.* 2:111-115.
- Boyd LB, Maynard MJ, Morgan-Lennell SK, Horton LB, Sugang R, Hamill RJ, Jimenez JR, Versalovic J, Steffen D, Zechiedrich L (2009). Relationship among Ciprofloxacin, Levofloxacin, and Norfloxacin MICs for Fluoroquinolone-Resistant *Escherichia coli* Clinical Isolates. *Antimicrob. Agents Chemother.* 53:220-234.
- Chitemerere TA, Mukanganyama S (2011). In vitro antibacterial activity of selected medicinal plants from Zimbabwe. *Afr. J. Plant Sci. Biotechnol.* 5:1-7.
- Cimanga K, De Bryune T, Pieters L, Tona L, Kambu K, Vanden Berghe D, Vlietinck AJ (1998). Antibacterial and antifungal activities of neocryptolepine, biscryptolepine and cryptoquindoline, alkaloids isolated from *Cryptolepis sanguinolenta*. *Phytomedicine* 5:209-214.
- Cowan MM (1999). Plant products as antimicrobial agents. *Clin. Microbiol. Rev.* 12:564-582.
- Das K, Tiwan RKS, Shrivastava DK (2010). Techniques for evaluation of medicinal plant products as antimicrobial agent: Current methods and future trends. *J. Med. Plants Res.* 4:104-111.
- Dash GK, Murthy PN (2011). Wound healing effects of *Ageratum conyzoides* Linn. *Int. J. Pharm. Bio. Sci.* 2:369-382.
- David O, Sanchez S, Gema J, NaCesa B, con Rivera I, Ramirez OD, Cumeros MCV, Garcia VMN (2009). Antimicrobial activity of medicinal plants from the Huautla Sierra biosphere reserve in Moreos Mexico. *Polibotanica Num.* 22:213-225.
- Daye NTK, Neuwman MJ, Smith EA, Tayma FSK (2008). Screening for antimicrobial activity of *Deferatum conyzoides* L.: A Pharmacological Approach. *Int. J. Pharmacol.* 5(2):14-18.
- Debajit K, Jayabrata S, Bharti S (2012). Ethno botanical antibacterial and antifungal study of three medicinal plants of Morigaon district of Assam, India. *Asian J. Pharm. Biol. Res.* 2:234-239.
- Djeussi DF, Noumedem JA, Seukep JA, Fankam AG, Voukeng IK, Tankeo SB, Nkuete AH, Kuste V (2013). Antibacterial activities of selected edible extracts against multidrug -resistant gram-negative bacteria. *Complement. Altern. Med.* 13:164.
- Djoukeng JB, Abou -Mansour E, Tabacchit B, Taponjon AL, Bouda H, Lontsi D (2005). Antibacterial triterpenes from *Syzygium guineense* (Myrtaceae). *J. Ethnopharmacol.* 101:283-286.
- Dugler B, Dugler G (2012). Antimicrobial activity of the leaves of *Ballota acetabulosa* on microorganisms isolated from urinary tract infections. *Turk. J. Pharm. Sci.* 9:257-262.
- Ebi GC, Ofoefule SJ (1997). Investigating into folkloric antimicrobial activities of *Landolphia owerience*. *Phytother. Res.* 11:149-155.
- Fabry W, Okemo PO, Musong R (1998). Antibacterial activity of East African medicinal plants. *J. Ethnopharmacol.* 60:79-81.
- Gamesan I, Krishnaraju I (1995). Antifungal properties of wild plants. *Adv. Plant Sci.* 8:194-196.
- Gupta PD, Babu R, Deshmukh PV (2013). Antibacterial activity of Indian medicinal plants against bacterial pathogens isolated from various skin ailments. *Asian J. Exp. Biol. Sci.* 4:336-344.
- Harborne JB (1998). *Phytochemical Methods. A guide to modern techniques of plant analysis.* Chapman & Hall, London.
- Hardman JG, Limbird LE (2001). *Goodman & Gilman's. The pharmacological basis of therapeutics*, 10th edition. Medical Publishing Division, New York. P 1163.
- Haruna MT, Anokwuru CP, Akeredolu AA, Akinsemolu AA, Alabi OA (2013). Antibacterial and antifungal activity of *Acalypha wilkesiana*. *Eur. J. Med. Plants* 3:52-64.
- Helberg I, Helberg D, Madau P, Mshingini KE, Mshiu EN, Samuelson G (1983). Inventory of plants used in traditional medicine in Tanzania. V. Plants of the family Dilleniaceae to Ophiliaceae. *J. Pharmacol.* 9:105-128.
- Islam MJ, Barua S, Das S, Khan N, Ahmed A (2008). Antibacterial activity of some indigenous medicinal plants. *J. Soil Nat.* 2:26-28.
- Israr F, Hassan F, Naqvi BS, Azhar I, Jabeen S, Hasan SM (2012). Report: Studies on antibacterial activity of some traditional medicinal plants used in folk medicine. *Pak. J. Pharm. Sci.* 25:669-674.
- Jagessar RC, Marsn A, Gomes O (2008). Selective antibacterial properties of *Phyllanthus acidus* leaf extract against *Candida albicans*, *Escherichia coli* and *Staphylococcus aureus* using Stokes Disc diffusion, Well diffusion Starch plate and a dilution method. *Nature Sci.* 6:24-38.
- Jamuna Bai A, Ravishankar RV, Pradeepa VS (2011). Evaluation of the antimicrobial activity of three medicinal plants of South India. *Malaysian J. Microbiol.* 7:14-18.
- Jekayinfa AO, George AO, Jayeoba KT (1997). *Acalypha wilkesiana*: preliminary in vitro microbiological and clinical trial on dermatitis. *Afr. J. Health Sci.* 4:39-42.
- Kambu K (1990). *Éléments de Phytothérapie Comparés. Plantes Médicinales Africaines.* Centre de Recherches Pédagogiques, Zaire.
- Kambu K, Tona L, Luki N, Cimanga K, Uvoya A (1990). *Activité antibactérienne d'extraits de plantes utilisées en préparation comme*

- antidiarrhéiques à Kinshasa, Zaïre. *Ann. Pharm. Fr.* 48:255-263.
- Kelmanson JE, Tager AAK, van Staden J (2000). Zulu medicinal plants used with antibacterial activity. *J. Ethnopharmacol.* 60:241-246.
- Kerharo J, Adam GA (1974). *Pharmacopée Sénégalaise Traditionnelle. Plantes médicinales et toxiques.* Edition Vigot Frères, Paris.
- Kondo S, Sattaponpan C, Phongpaicht S, Srijan A, Itharat A (2010). Antibacterial activity of Thai medicinal plants *Pikutbenjakul*. *J. Med. Assoc. Thai.* 7:331-135.
- Kuphumla Z (2014). *Sygium guineense* (Wild.) DC. Seedroom, Kirstenbosh National Botanical Gardens. <http://www.plantzafrica.com/plantqrs/syzygiumguin.htm>
- Lim SW, Loh HS, Ting KN, Bradshaw TD, Zeenzthul NA (2013). *Acalpha wilkesiana* ethyl acetate extract enhances the in vitro cytotoxic effects of -tocopherol in human brain and lung cancer cells. *Int. J. Biosci. Biochem. Bioinf.* 3:335-340.
- Lotifpour F, Nazemiyeh N, Fathi-Azad F, Garacci N, Arami S, Talat S, Sadegjsour F, Hasanporia B (2008). Evaluation of the antibacterial activity of some medicinal plants from North-West Iran. *Ir. J. Basic Med. Sci.* 11:80-85.
- Malesh B, Satish S (2008). Antimicrobial activity of some important medicinal plants against plant and human pathogens. *World J. Agric. Sci.* 4:839-843.
- Maroyi A (2008). *Syzygium* (Wild.) DC. In: Louppe D, Oteng-Amoako AA, Brik M (Eds.), *Protas* 7(1): Timbers/Bois d'oeuvre 1 [CD]. PROTA, Wageningen. Netherlands. Available on http://database.prota.org/PROTAhtml/Syzygium%20guineense_En.htm
- Mendonça Filho RR (2006). To modern phytomedicine turning medicinal plants into drugs. Bioactive phytochemicals. In: Ahmadd J, Agil P, Owans M (Eds.), *New approaches in the Phytosciences.* Wiley-VCH, Verlag, GmbI & Co. KGaA, Weinheim. pp. 1-24.
- Muhamed Mubarak H, Doss A, Dhanabalan R, Venkataswamy R (2011). In-vitro antimicrobial effects of some selected plants against bovine mastitis pathogens. *Hygeia. J. Drugs Med.* 3:71-75.
- Mustafa MR, Mahmood AA, Sidik K, Noor SM (2005). Evaluation of wound healing potential of *Ageratum* leaf extract in combination with honey in rats as a minor model. *Int. J. Mol. Adv. Sci.* 1:406-410.
- Neeraj T, Madivi A (2011). Studies on antibacterial activity of methanol plant extracts. *Int. J. Phytomed.* 3:549-556.
- Neuwinger HD (2000). *African Traditional Medicine. A Dictionary of Plant Uses and Application.* Medipharm Scientific Publishers, Stuttgart. pp. 421-422.
- Okwori A, Dina C, Jumaid S, Okeke I, Adetunji J, Olabode A (2013). Antibacterial activities of *Ageratum conyzoides* on selected bacterial pathogens. *Int. J. Microbiol.* 4(1):1-4.
- Oliver-Bever B (1986). *Medicinal plants in tropical West Africa.* Cambridge, London.
- Osho A, Adetunji T (2011). Antibacterial activity of the essential oil of *Ageratum conyzoides* L. *Asian J. Sci. Technol.* 2:1-5.
- Oskay M, Oskay D, Kalyoncu F (2009). Activity of some plant extracts against multi-drug resistant human pathogens. *Iranian J. Pharm. Res.* 8:293-300.
- Rabe T, van Saden I (1997). Antibacterial activity of South African plants used for medicinal purposes. *J. Ethnopharmacol.* 56:81-87.
- Raghavendra R, Mahadevan GD (2011). *In vitro* antimicrobial activity of various plant latex against resistant human pathogens. *Int. J. Pharm. Sci.* 3:70-72.
- Rios JL, Recio MC (2005). Medicinal plants and antibacterial activity. *J. Ethnopharmacol.* 100:80-84.
- Royal Botanic Gardens, Kew (2014). Entry for *Acalypha wilkesiana* Müll.-Arg. [family Euphorbiaceae] Advanced Search. http://plants.jstor.org/upwta/2_56
- Sadeghi-Nejad B, Azish M (2013). In vitro antibacterial and antifungal effect of some medicinal plants. *Afr. J. Microbiol. Res.* 7, 3802-3806.
- Sakunpak A, Panichayupakaranant P (2012). Antibacterial activity of Thai edible plants against gastrointestinal pathogenic bacteria and isolation of new broad spectrum antibacterial polyisoprenylated benzophenone, chamuangone. *Food Chem.* 130:826-831.
- Samy RP, Ignacinuthu S (2000). Antibacterial activity of some folklore medicinal plants used by tribals in Western Ghat of India. *J. Ethnopharmacol.* 69:63-71.
- Sharmeen R, Hossain N, Rrahman M, Foysal J, Miah F (2012). In-vitro antibacterial activity of herbal aqueous extract against multi-drug resistant *Klebsiella* sp. isolated from human clinical samples. *Int. Curr. Pharm. J.* 1:133-137.
- Shinkafi SA, Dauda H (2013). Antibacterial activity of *Allium cepa* (Onion) on some pathogenic bacteria associated with ocular infections. *Sch. J. App. Med. Sci.* 1:147-154.
- Singla RK, Jaiswal N, Bhat GV, Jagani H (2011). Antioxydant and antimicrobial activities of *Coco nucifera* Linn. (Arecaceae) endocarp extracts. *Indo Global J. Pharm. Sci.* 1:354-361.
- Tabassum H, Naiman A, Al-Noura M, Al-Jameil A, Khan F (2013). Evaluation of antibacterial potential of selected plant extracts on bacterial pathogens isolated from urinary tract infections. *Int. J. Curr. Microbiol. App. Sci.* 2:353-368.
- Tona L, Kambu K, Mesia K, Cimanga K, Apers S, De Bruyne T, Pieters L, Totté J, Vlietinck AJ (1999). Biological screening of traditional preparations from medicinal plants used as antidiarrhoeal in Kinshasa, Congo. *Phytomedicine* 61:56-66.
- Vanden Berghe D, Vlietinck AJ (1991). Screening for antibacterial and antiviral agents. In: Hostettman K (Ed.), *Plant Biochemistry, Vol. 6. Assay for Bioactivity.* Academic Press, London. pp. 47-70.
- Vlietinck AJ, Van Hoof I, Totté J, Lasure A, Vanden Berghe D, Rwangabo PC, Mvukiyumwanu J (1995). Screening of hundred Rwandese medicinal plants for anti-microbial and antiviral properties. *J. Ethnopharmacol.* 46:31-37.
- Wendakoon C, Calderon P (2012). Evaluation of selected medicinal plants extracted in different ethanol concentration for antibacterial activity against human pathogens. *J. Med. Active Plants* 1:60-68.
- Wheels M (2008). *Principles of modern microbiology.* Jones and Bartlett Publishers, Inc, Boston.
- World Health Organization (WHO) (2002). *Traditional medicine Growing Needs and Potential-WHO Polley Perspectives on Medicine.* No 002, May, World Health Organisation, Geneva, Switzerland.
- Yamamoto IA, Sildera JC, Emim JA, Godinho RO, Souccar C, Lapa AJ (1991). Pharmacological screening of *Ageratum conyzoides* L. *Mem. Inst. Oswaldo Cruz.* 86(Suppl 2):145-147.