

Full Length Research Paper

# Iron content of the formulated East African indigenous vegetable recipes

Florence O. Habwe<sup>1\*</sup>, Mary K. Walingo<sup>1</sup>, Mary O. Abukutsa-Onyango<sup>2</sup> and Mel O. Oluoch<sup>3</sup>

<sup>1</sup>School of Public Health, Maseno University, Box 333–40105, Maseno, Kenya.

<sup>2</sup>Horticulture Department, Jomo Kenyatta University of Agriculture and Technology, Box 62000, 00200, Nairobi, Kenya

<sup>3</sup>World Vegetable Center -Regional Center for Africa, Box 10, Duluti, Arusha, Tanzania.

Accepted 02 November, 2019

Nutritional iron deficiency is the common cause of anaemia in developing world. However, Africa is endowed with African indigenous vegetables (AIVs) rich in micronutrients. Surveys within East Africa relate low AIV consumption to limited information on recipe preparation. The aim of this research is to determine iron content of various formulated indigenous vegetable recipes. Experimental research involved four randomly selected AIVs; African Nightshade (*Solanum scabrum*), Vegetable Amaranth (*Amaranthus blitum*), Slenderleaf (*Crotalaria ochroleuca*) and Cowpea (*Vigna unguiculata*). Four single and six vegetable combinations were boiled for ten minutes with and without traditional salt, then fried for five minutes using onions and tomatoes, giving rise to twenty recipes. Iron content was evaluated using Atomic Absorption Spectroscopy. Iron content of raw indigenous vegetables was significantly ( $P < 0.05$ ) lower than boiled and fried AIVs. Fried AIVs had significantly ( $P < 0.05$ ) higher iron content compared to the boiled AIVs. Mean iron content of AIVs fried with lye was insignificantly ( $P > 0.05$ ) lower compared to those fried without lye. Fried AIVs should be recommended in areas with high dietary iron deficiency; this could help alleviate anaemia.

**Key words:** Iron content, recipes, lye, African indigenous vegetables.

## INTRODUCTION

Indigenous vegetables are vegetable species native to or originating from a particular region or environment, are acceptable and used in the diet of rural and urban people

\*Corresponding author. E-mail: [habwerence@yahoo.com](mailto:habwerence@yahoo.com) or [habwerence@googlemail.com](mailto:habwerence@googlemail.com). Tel: +254 713 741144.

**Abbreviations:** AIVs, African Indigenous Vegetables; AVRDC, Asian Vegetables Research and Development Center; AAS, Atomic Absorption Spectroscopy; ACC, Administrative Committee on Coordination; FORMAT, Forum for Organic Resource Management and Agricultural Technologies; IVs, Indigenous Vegetables; ICRAF, International Center for Research in Agroforestry, IPGRI, International Plant Genetic Resources Institute, IDA, Iron Deficiency Anaemia; KENRIK, Kenya Resource Center for Indigenous Knowledge; NEVEPA, Network Vegetable Production in Africa; NNMK, National Museums of Kenya; SCN, Sub-Committee on Nutrition; UNICEF, United Nations Children's Fund; VICRES, Lake Victoria Research Initiative; WAC, World Agroforestry Center; WHO, World Health Organization; WVC, World Vegetable Center.

(Maundu, 1997; FAO, 1988). African indigenous vegetables can play a significant role in addressing three major factors of low income, malnutrition and loss of biodiversity. The diversity of these African indigenous vegetables forms the basis of food and nutrient diversity (FAO, 2003). These vegetables are still valuable sources of nutrients thus contribute substantially to vitamin and mineral intake (Waudu et al., 2007; Farm Africa, 2006; Abukutsa-Onyango, 2003). They contribute essential components of the meal by contributing micronutrients like iron which are usually in short supply in daily diets, especially where animal protein is unaffordable (Mosh and Gaga, 1999). Some of these vegetables are gathered when in season, grown in home gardens or intercropped with staples; and eventually find their way to the urban markets (Mnzava, 1995).

AIVs can be used as cash crops in peri-urban systems, source of vegetables for daily sustenance in home gardens, source of new crops and as source of variation for diversification of production systems and diet. While most AIVs are nutritious if well cooked, a survey carried out by Waudu et al. (2005) in wetlands of Lake Victoria

region and that of Waudu et al. (2007) in urban and peri-urban settings of Nairobi Kenya, shows low intakes of AIVs by women and children due to lack of knowledge on preparation and cooking procedures. The later survey shows that respondents used AIV cooking procedures that could lead to a decrease of the nutritive value of cooked vegetables. It is therefore no wonder that micro-nutrient deficiency mainly vitamins and minerals particularly iron is a serious problem in sub-Saharan Africa affecting about a third of the population with far reaching effects (UNICEF, 2004).

Survey results by Waudu et al. (2007) indicate the need to discourage chopping vegetables before washing, repeated boiling and adding of sodium bicarbonate to vegetables while cooking. However, there has been very little information on recipes and their nutrient content which has been in high demand (Shiundu and Oniang'o, 2007). This shows the urgency need of information and knowledge on AIV processing and preparation alongside their nutrient content after cooking; as agreed by Padma, 2005, who stated that it is not enough to develop recipes but their nutrient content is important in the effort of improving nutrition security. The objective of this study was to develop East African indigenous vegetable recipes and evaluate their iron content.

## **MATERIALS AND METHODS**

### **Study site**

The study was carried out at Maseno University, Kenya. Long-term average rainfall in Maseno is 2074 mm per annum and its distribution is bimodal with peaks in March/April and September/October (Oseko, 2007). Soils are mainly dominated by vertisols, with a fairly acidic pH in water of 4.5 to 6.5 (Otieno et al., 1993; GOK, 2002; and Oseko, 2007). The soils are also deep, very deficient in potassium (P) and Nitrogen (N) and have moderate potassium (P) fixation (FAO,1997, cited by Oseko, 2007). Mean annual day temperature is 20°C with the average maximum daily temperature not exceeding 31°C and the average minimum night temperature above 15°C (Otieno et al., 1993; GOK, 2002; Oseko, 2007).

### **Source of East African indigenous vegetables**

African indigenous vegetables used in the experiment were planted at the Maseno University Botanic garden. Land was prepared by ploughing and harrowing to a fine tilth. A plots of 5 × 5 m were demarcated and poultry manure was mixed with the soil in the demarcated plot at a rate of 5 tons per ha. Seeds of each selected AIV were mixed with the soil at a rate of 1:10 and drilled in the respective plots at a spacing of 30 cm. After two weeks, thinning was done to leave an inter-row spacing of 15 cm for all. All other agronomic practices were done to ensure optimum growing conditions. Harvesting was done at four weeks after seedling emergency. This resulted to AIVs with similar harvest age and environmental exposure.

**Formulation of East African indigenous vegetable recipes** After harvesting the AIVs, immediately destalking was done to se-

parate vegetable leaves from the stems in order to minimize nutrient loss. Required amounts of AIVs were then weighed and various vegetable categories of single vegetables and vegetable combinations were made. This resulted into four single vegetables and 6 vegetable combinations that were washed to remove dirt and then boiled with and without traditional salt for ten minutes over moderate heat at 40°C. After boiling, vegetables were put aside and vegetable oil was put on a pan on fire to heat, diced onions were fried till golden brown, chopped tomatoes added and cooked till soft while stirring to improve taste and aroma. The boiled vegetables were then added, stirred and common salt was put to taste. This was simmered for two minutes, removed from fire and served.

The traditional salt (lye) was prepared by drying the pods of green beans after removing the mature seeds, the dry pods were then burnt over a hot dry pan and the ash collected after complete burning. The ash was put in a container whose bottom had small holes and water poured in it to pass through the ash into another container underneath. The residue is what is known as traditional salt or lye.

### **Sample preparation for elemental analysis**

After cooking, samples were cooled and immediately kept in the fridge at 4°C to prevent distortion of the cooking ingredients. Samples were oven-dried at 60°C for 12 h then crushed into fine powder using a mill (QCG System LLC Model 4E). The powdered samples were sieved through 125 µm apertures BS 410. 0.5 g of each sample was weighed and placed in a kjeldahl tube in 20 mls of aqua regia acid (5 ml HCl + 5 ml Nitric acid) (Apha, 1985). The tubes were then heated to boiling for 2 h at 96°C until the resulting solution was clear and the heating was continued for 30 more minutes. The digested samples were left to cool overnight and the contents were transferred into a 50 ml volumetric flask and made up to the mark with distilled water. The samples were then filtered through a Whatman filter paper No.1 and finally transferred to 100 ml polypropylene bottles, ready for elemental analysis.

### **Evaluation of iron content of selected East African indigenous vegetables**

Elemental analysis was done in three states: on raw AIVs, after boiling and after frying the AIVs. All the material was taken for analysis as there was no liquid left behind after cooking. Atomic Absorption Spectroscopy (AAS) was used where standards of each element (1000 parts per million) were diluted with 10% HCl. This was then aspirated directly using a Narian AAS Model at Mines and Geology department, Ministry of Natural Resources, Nairobi. Kenya (Gerge, 1984).

### **Data analysis**

Data obtained were analyzed using ANOVA, descriptive and inferential statistics. Iron content data were subjected to independent and paired sample t- test to determine whether the treatments' effects were significant at 5% level of significance.

## **RESULTS**

### **The formulated African indigenous vegetable recipes**

Twenty East African indigenous vegetable recipes were formulated in the study. All these vegetable recipes

**Table 1.** Summary of the Fried Vegetable Recipes.

| Recipe                                 | Quantity of ingredients for each recipe     |
|--|---|
| Nightshade with lye                    | ½ kg AIVs for single vegetable recipes      |
| Nightshade without lye                 | ¼ kg AIVs for vegetable combination recipes |
| Cowpea with lye                        | ¼ liter of water                            |
| Cowpea without lye                     | 1 medium diced onion                        |
| Slenderleaf with lye                   | 1 tbsp vegetable oil                        |
| Slenderleaf without lye                | 2 medium tomatoes                           |
| Amaranth with lye                      | 2 tbsp table salt                           |
| Amaranth without lye                   | 3 tbsp traditional salt (lye)               |
| Nightshade and Amaranth with lye       |   |
| Nightshade and Amaranth without lye    |   |
| Nightshade and Slenderleaf with lye    |   |
| Nightshade and Slenderleaf without lye |   |
| Nightshade and Cowpea with lye         |   |
| Nightshade and Cowpea without lye      |   |
| Amaranth and Slenderleaf with lye      |   |
| Amaranth and Slenderleaf without lye   |   |
| Amaranth and Cowpea with lye           |   |
| Amaranth and Cowpea without lye        |   |
| Slenderleaf and Cowpea with lye        |   |
| Slenderleaf and Cowpea without lye     |   |

**Table 2.** Iron content (mg/g) of selected AIVs under different treatments.

| AIVs                       | Raw   | Boiled with lye | Boiled no lye      | Fried with lye | Fried no lye | Average              |
|----------------------------|-------|-----------------|--------------------|----------------|--------------|----------------------|
| Nightshade                 | 17.3  | 12.4            | 11.5               | 281.2          | 401.6        | 144.8 <sup>ab</sup>  |
| Cowpea                     | 24.1  | 20.6            | 15.3               | 16.7           | 1208         | 256.94 <sup>ab</sup> |
| Slenderleaf                | 14.7  | 10.7            | 6.4                | 110            | 5.8          | 29.52 <sup>b</sup>   |
| Amaranth                   | 19.4  | 95.2            | 12.4               | 108            | 5.3          | 48.06 <sup>b</sup>   |
| Nightshade and Amaranth    | 20.5  | 11.5            | 11.5               | 9.1            | 42.3         | 18.98 <sup>b</sup>   |
| Nightshade and Slenderleaf | 13.2  | 12.2            | 13.9               | 5.1            | 8.2          | 10.52 <sup>b</sup>   |
| Nightshade and Cowpea      | 23.3  | 69.2            | 200.4              | 381.4          | 6.6          | 136.18 <sup>ab</sup> |
| Amaranth and Slenderleaf   | 12.5  | 23.6            | 557.5              | 859.2          | 859.2        | 469.6 <sup>a</sup>   |
| Amaranth and Cowpea        | 20.1  | 69.2            | 16.6               | 7.2            | 303          | 83.22 <sup>b</sup>   |
| Slenderleaf and Cowpea     | 14.7  | 26.9            | 9.1                | 10.1           | 8.1          | 13.78 <sup>b</sup>   |
| Average                    | 17.98 | 35.15           | 85.46              | 178.8          | 284.81       | 120.44               |
| Significance Level         |       |                 |                    |                |              | 0.05                 |
| LSD                        |       |                 |                    |                |              | 348.28               |
| Interaction                |       |                 | Cooking Method*Lye |                |              | ns                   |
|                            |       |                 | Cooking Method*AIV |                |              | ns                   |

NB: Traditional Salt (Lye) = 7.2 mg/g.


(Table 1) were prepared using the same methodology.

### Iron content of commonly consumed African indigenous vegetables

Iron content of selected AIVs and its combinations are indicated on Table 2 and it is demonstrated that cooking has released more extractable iron in African indigenous

vegetables. There were no significant interactions between cooking method with AIVs (Table 2); however a combination of amaranth and slenderleaf had significantly higher iron solubility. Apart from nightshade with amaranth, nightshade with slenderleaf and amaranth with cowpea, all other AIVs had their iron solubility enhanced after frying (Table 2).

Whether the vegetables were cooked as single vegeta-



**Figure 1.** General effect of cooking and cooking method on iron content of AIVs.

bles or as a combination of two vegetables did not have an effect on their iron solubility (Table 2), some vegetables had higher iron content as single vegetables while others had higher content when combined with others. For example, a combination of amaranth and slenderleaf boiled without lye, fried with lye and fried without lye had the highest extractable iron of 557, 859.2 and 859.2 mg/g, respectively compared to amaranth and slenderleaf alone (Table 2).

Raw African indigenous vegetables had the least mean iron content compared to the boiled and the fried AIVs. Recipes prepared without traditional salt, recorded higher mean iron content for fried (without lye 284.8 + 427, with traditional salt 272.7 + 382.6) compared to mean iron content for boiled (without traditional salt 85 + 176, with traditional salt 30.7 + 28.4). Fried AIVs had higher mean iron content compared to the boiled (Figure 1). Paired sample t-test was applied and the results indicated that fried AIVs had significantly higher iron content compared to the boiled AIVs ( $P < 0.05$ ). The difference in mean iron content between AIVs fried with lye and those fried without lye was insignificant ( $P > 0.05$ ).

## DISCUSSION

### Formulation of African indigenous vegetable recipes

Lack of variety in recipe formulation has in the past been an obstacle towards AIV consumption and marketability according to Abukutsa-Onyango et al. (2006) and Shiundu and Oniang'o (2007). Therefore availability of AIV recipes could pave way for increased AIV consumption, ensure availability throughout the year and improve marketability.

### Iron content of commonly consumed AIVs

Results indicate that cooking increase iron in AIVs; this concur with the experimental findings of Imungi and Poter

(1983)( as cited by Abukutsa-Onyango, 2003) which showed increase in iron content of some African indigenous vegetables after cooking. This is because the heat exposes solubility of iron thus aids its bioavailability. The recommended daily intake (RDI) for iron content is between 10 to 8 mg (NIN, 1992; Abukutsa-Onyango, 2003); (Table 2) most formulated recipes could supply the recommended daily intake.

A study carried out by Weinberger and Msuya (2004) indicated that raw amaranth and nightshade contain iron content of 37.05 and 8.90 mg/g, respectively; this slightly varies with the results of this study, which indicate iron contents of 19.4 and 17.3 mg/g respectively. The possible explanation for the slight differences in iron contents could be due to the differences in type of vegetable and place (plots) where they were obtained (Weinberger and Msuya, 2004).

Combining vegetables during preparation have different effects on different vegetables (Table 2) in terms of their iron solubility and this could be attributed to different nutrient-nutrient interactions between different vegetables. Frying significantly ( $P < 0.05$ ) increased iron solubility (Table 2); this could be attributed to the increased cooking time and also more iron could have been generated from the onions and tomatoes used during frying which are known to contain 1.48 and 1.99 mg/100g of iron, respectively (Sehmi, 1993) and vitamin A in the oil that increase iron solubility.

## ACKNOWLEDGEMENTS

The authors are grateful for the research grants provided by the project on the Promotion of Neglected Indigenous Vegetables Crops for Nutritional Health in East and Southern Africa Phase 2 (ProNIVA2), which was awarded by AVRDC-RCA (The World Vegetable Centre – Regional Centre for Africa) . The African Indigenous Vegetable seeds used in this research were provided by VicRes AIVs Project funded by SIDA through the Lake Victoria Research Initiative – VicRes.

## REFERENCES

- Abukutsa-Onyango MO (2003). Unexploited Potential of Indigenous African Vegetables in Western Kenya. Department of Botany and Horticulture. Maseno University. Kenya. 4: 103-122.
- Abukutsa-Onyango MO, Tushaboomwe K, Onyango JC, Macha SE (2006). Improved Community Landuse for Sustainable Production and Utilization of African Indigenous Vegetables in the Lake Victoria Region. In: Proceedings of the Fifth Workshop on Sustainable Horticultural Production in the Tropics 23rd – 26th November 2005, ARC, Egerton University, Njoro: Egerton University. pp 167-179.
- Apha A (1985). Standard Methods for the Examination of Water and Wastewater. 16<sup>th</sup> edition. WPCF.
- FAO (1997). Preventing Micronutrient Malnutrition: A guide to food-based approaches – A Manual for Policy Makers and Programme Planners. Rome: FAO.
- FAO (1988). Traditional Food Plants. Food and Nutrition Paper 42/1 FAO. Rome. Italy.
- Farm Africa (2006). Research Information Ltd. <http://www.appropriate-technology.org>. Accessed Feb 26<sup>th</sup> 2008.
- Gerge C (1984). Analysis of foods and Beverages Modern Technologies. Accademic Press. INCI.
- Imungi JK, Porter NN (1983). Nutrient Contents of Raw and Cooked Cowpea Leaves. J. Food Sci. 48: 1252-1254.
- Kenya Government (2002). Effective management for sustainable economic growth and poverty reduction. Nairobi: Ministry of Finance and Planning Government of Kenya.
- Maundu PM (1997). The status of traditional vegetable utilization in Kenya. In: Proceedings of the IPGRI International Workshop on Genetic Resources of Traditional Vegetables in Africa, Conservation and Use. ICRAF-HQ, Nairobi: IPGRI.
- Mnzava NA (1995). Traditional Vegetables in Tanzania. Paper presented to the 1<sup>st</sup> National Research Planning Workshop, Arusha. Tanzania.
- Mosha TC, Gaga HE (1999). Nutritive value and effect of blanching on trypsin and chymotrypsin inhibitor activities of selected leafy vegetables. J. Plant Foods Hum. Nutr. 54: 271-283.
- NIN (1992). Dietary Guidelines for Indians – a manual. Hyderabad, Indian Council of Medical Research. NIN.
- Oseko JK (2007). Performance of African Indigenous Vegetables under Sole and Intercropping Systems: A case of African Kale (*Brassica carinata*). An unpublished Msc thesis of Maseno University. Kenya.
- Otieno HJO, Amadalo B, Gathumbi S (1993). Afrena Project Maseno Kenya. Maseno: ICRAF.
- Padma TV (2005). Beating World Hunger. The return of 'Neglected' Crops. <http://www.scidev.net>.
- Sehmi JR (1993). National Food Composition Tables and the Planning of Satisfactory Diets in Kenya. Nairobi: Government Printer.
- Shiundu KM, Oniang'o R (2007). Marketing African Leafy Vegetables, Challenges and Opportunities in the Kenyan Context. J. Food Agric. Nutr. Dev. 17: 4.
- UNICEF (2004). Report on Vitamin and Mineral Deficiency in Sub-Saharan Africa.
- Waudu J, Tuitoek P, Msoya J, Kikafunds J (2005). Food Consumption Patterns and Nutrient Intakes by Women and Children Under-five in the Wet lands of Lake Victoria Basin. Research Report for VICRES.
- Waudu J, Kimeywe J, Mbithe D, Maundu P (2007). Utilization and Medical Value of Indigenous Leafy Vegetables Consumed in Urban and Peri-urban Nairobi. Provide title of paper African journal of food and agriculture nutrition and development. ISSN 1684-5374. 7:4.
- Weinberger K, Msuya J (2004). Indigenous Vegetables in Tanzania – Significance and Prospects. Shanhu, Taiwan: AVRDC – The World Vegetable Center, Technical Bulletin No 31, AVRDC Publication 04-600. p. 70